

## Metabolic profiling of adiponectin levels in adults: Mendelian randomization analysis

Borges, Maria Carolina<sup>1, 2,3</sup>, carolina.borges.mcb@gmail.com

Barros, Aluísio JD<sup>1</sup>, abarros.epi@gmail.com

Ferreira, Diana L Santos<sup>2,3</sup>, diana.santosferreira@bristol.ac.uk

Casas, Juan Pablo<sup>4</sup>, jp.casas@ucl.ac.uk

Horta, Bernardo Lessa<sup>1</sup>, blhorta@gmail.com

Kivimaki, Mika<sup>5</sup>, m.kivimaki@ucl.ac.uk

Kumari, Meena<sup>5,6</sup>, mkumari@essex.ac.uk

Usha, Menon<sup>7</sup>, u.menon@ucl.ac.uk

Gaunt, Tom R<sup>2,3</sup>, tom.gaunt@bristol.ac.uk

Ben-Shlomo, Yoav<sup>3</sup>, y.ben-shlomo@bristol.ac.uk

Freitas, Deise F<sup>1</sup>, ise.freitas07@hotmail.com

Oliveira, Isabel O<sup>1,8</sup>, isabel.ufpel@gmail.com

Gentry-Maharaj, Aleksandra<sup>7</sup>, a.gentry-maharaj@ucl.ac.uk

Fourkala, Evangelia<sup>7</sup>, rania4kala@yahoo.co.uk

Lawlor, Debbie A<sup>2,3,\*</sup>, D.A.Lawlor@bristol.ac.uk

Hingorani, Aroon D<sup>4,9,\*</sup>, a.hingorani@ucl.ac.uk

\* Joint last authors

<sup>1</sup> Post-Graduate Program in Epidemiology, Federal University of Pelotas, Pelotas, Brazil.

<sup>2</sup> MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK

<sup>3</sup> School of Social and Community Medicine, University of Bristol, Bristol, UK

<sup>4</sup> Farr Institute of Health Informatics, University College London, London, UK

<sup>5</sup> Department of Epidemiology and Public Health, University College London, London, UK

<sup>6</sup> Institute for Social and Economic Research, University of Essex, Essex, UK

<sup>7</sup> Department of Women's Cancer, Institute for Women's Health, Faculty of Population Health Sciences, University College London, London, UK

<sup>8</sup> Department of Physiology and Pharmacology, Institute of Biology, Federal University of Pelotas, Pelotas, Brazil

<sup>9</sup> Institute of Cardiovascular Science, University College London, London, UK

Corresponding author: Borges MC. Rua Marechal Deodoro, 1160 - 3° Piso, Centro, Pelotas, RS, Brazil. Zipcode: 96020-220. Phone: +55 (53) 3284 - 1300. Email: carolina.borges.mcb@gmail.com

## ABSTRACT

**Background:** Adiponectin, a circulating adipocyte-derived protein has insulin-sensitizing, anti-inflammatory, anti-atherogenic, and cardiomyocyte-protective properties in animal models. However, the systemic effects of adiponectin in humans are unknown.

**Objectives:** Our aims were to define the metabolic profile associated with higher blood adiponectin concentration and investigate whether variation in adiponectin concentration affects the systemic metabolic profile.

**Methods:** We applied multivariable regression in up to 5,906 adults and Mendelian randomization (using *cis*-acting genetic variants in the vicinity of the adiponectin gene as instrumental variables) for analysing the causal effect of adiponectin in the metabolic profile of up to 38,058 adults. Participants were largely European from six longitudinal studies and one genome-wide association consortium.

**Results:** In the multivariable regression analyses, higher circulating adiponectin was associated with higher HDL lipids and lower VLDL lipids, glucose levels, branched-chain amino acids, and inflammatory markers. However, these findings were not supported by Mendelian randomization analyses for most metabolites. Findings were consistent between sexes and after excluding high risk groups (defined by age and occurrence of previous cardiovascular event) and one study with admixed population.

**Conclusion:** Our findings indicate that blood adiponectin concentration is more likely to be an epiphenomenon in the context of metabolic disease than a key determinant.

## INTRODUCTION

The recognition that adipose tissue is an endocrine organ raised new prospects for discovering adipose-derived products that could be valuable drug targets for the treatment and prevention of cardio-metabolic diseases. In this context, adiponectin, a 30KDa protein largely produced by mature adipocytes, has been attracting widespread attention due to insulin-sensitizing, anti-inflammatory, anti-atherogenic, and cardiomyocyte-protective properties demonstrated in animal models (1).

However, human studies have yielded a far more complicated picture. Unlike most other adipokines, circulating adiponectin concentration is higher with lower adiposity (2). In prospective observational studies in humans using multivariable regression, higher circulating adiponectin is associated with lower risk of type 2 diabetes (3), hepatic dysfunction (4), and metabolic syndrome (5), but higher mortality in patients with kidney disease, heart failure, previous cardiovascular disease or general elderly cohorts (6-9); this different direction of effect between risk of incident disease and mortality among high risk groups has been called “the adiponectin paradox” (10).

Given the complex metabolic derangements that might participate in and compensatory changes that might occur in response to human diseases, the association between adiponectin concentration and cardio-metabolic biomarkers and disease end-points might be explained by reverse causality (where disease status could alter adiponectin concentration) or residual confounding (where adiponectin could be a marker of another causal factor, such as adiposity or insulin resistance) (11). Classical multivariable regression studies cannot distinguish causal from non-causal associations, and randomized controlled trials (RCTs) specifically targeting adiponectin are not possible in the absence of a specific therapeutic targeting adiponectin concentration or function.

Mendelian randomization uses genetic variants (mostly single nucleotide polymorphisms (SNPs)) that are robustly related to the risk factor of interest as tools to assess its role in causing disease (12). The random allocation of parental alleles at meiosis should theoretically reduce confounding in genetic association studies and this has been shown to be the case (13); the unidirectional flow of biological information from genetic variant to phenotypes avoids reverse causality. Mendelian

randomization has been used in clinical research to investigate potential etiological mechanisms, such as the causal effects of low density lipoprotein cholesterol (LDL-c) (14), systolic blood pressure (SBP) (14) and C reactive protein (15) on coronary heart disease (CHD), validate and prioritize novel drug targets, such as interleukin-6 receptor (16), and increase understanding of current therapies, for example statins (17).

Previous Mendelian randomization studies indicate that circulating adiponectin is a consequence of low insulin sensitivity (18), but whether adiponectin concentration is also a cause of insulin sensitivity is uncertain (18-20). Using Mendelian randomization in a study of 63,746 CHD cases and 130,681 controls we have recently shown that adiponectin may not be causally related to CHD (21). Whilst multivariable analyses show higher adiponectin concentration is associated with lower glycated haemoglobin, insulin, triglycerides and higher high density lipoprotein-cholesterol (HDL-c), using Mendelian randomization, we found little evidence these were causal (21). Whether adiponectin is associated with systemic metabolic profile, and, if it is, what aspects of these associations are causal is unknown. A broader interrogation of the metabolic effects of adiponectin through high-throughput profiling of metabolic status could provide valuable insights into whether adiponectin is a non-causal biomarker or causally important in the pathophysiology of some human diseases (22).

We combined genotype, adiponectin and metabolomics profile data from six longitudinal studies and one genome-wide association consortium with the aim of (i) defining the metabolic effects of blood adiponectin concentration and (ii) investigating whether variation in adiponectin concentration is causally related to the systemic metabolic profile.

## METHODS

### *Study Populations*

The metabolic profile associated with blood adiponectin concentration was examined from seven data sources: the 1982 Pelotas Birth Cohort (PEL82), including adults aged 30 years old born in the city of Pelotas, Brazil, in 1982 (23, 24); the British Women's Heart and Health Study (BWHHS), including UK women aged 60-79 years old at recruitment in 2000 (25); the Whitehall II Study (WHII), including UK government workers aged 45-69 years at phase 5 clinical assessment in 1997-1999 (26); the Caerphilly Prospective Study (CaPS), including men aged 52-72 years at phase III in 1989-1993 (27); a case-control study nested in The United Kingdom Collaborative Trial of Ovarian Cancer Screening (UKCTOCS), including UK postmenopausal women aged 50-74 years at recruitment in 2001-2005 (28); the cohort of mothers from The Avon Longitudinal Study of Children and Parents (ALSPAC-M), including UK women aged 34-63 years old at clinical assessment in 2009-2011 (29); and a metabolomics genome-wide association consortium (Metabolomics consortium), including European adults with mean age of 45 years old from 14 cohorts (30). Individual level data was available to investigators from PEL82, BWHHS, WHII, CaPS, UKCTOCS and ALSPAC-M and summary level data is publicly available from the Metabolomics consortium (URL: [http://www.computationalmedicine.fi/data/NMR\\_GWAS/](http://www.computationalmedicine.fi/data/NMR_GWAS/)). All study participants provided written informed consent, and study protocols were approved by the local ethics committees (ethical approval for ALSPAC was also obtained from the ALSPAC Ethics and Law Committee). Studies' characteristics are summarized on **Table 1**. We examined (possibly causal) associations of adiponectin with systemic metabolic profiles using two approaches – conventional multivariable regression and Mendelian randomization analyses. Studies must have both adiponectin and measures of some of the outcomes (but do not need genetic data) to contribute to multivariable regression analyses, and must have relevant genetic variants and outcomes (but do not need adiponectin concentration data) to contribute to Mendelian randomization analyses. **Figure 1** shows how the different data sources contributed to the two approaches.

### *Metabolite Quantification*

A high-throughput serum nuclear magnetic resonance (NMR) spectroscopy platform was utilized to quantify up to 150 metabolic measures and 83 derived measures (ratios) in each study. This NMR platform has been used in several other studies (22, 31, 32) and methodological details have been described elsewhere (33, 34). 66 out of 150 metabolic measures were selected for this study aimed at broadly representing the systemic metabolite profile, as previously reported by Wurtz et al (35), including: lipoprotein traits (lipid content, particle size, and apolipoproteins), free fatty acids, amino acids, glycolysis-related metabolites, ketone bodies, fluid balance (albumin and creatinine), and inflammatory markers (glycoprotein acetyls). The remaining 84 metabolic measures from the NMR platform are related to other lipid fractions (esterified and free cholesterol, total cholesterol, triglycerides, and phospholipids) and particle concentration from 14 lipoprotein subclasses and are not presented in this study. Instead, we present the total lipid content of each of the 14 lipoprotein subclasses, which is highly correlated to their respective lipid fractions and particle concentration and comprehensively represents the plasma lipid partitioning across lipoproteins. Eight additional measures, not obtained from the NMR platform, were also included: C-reactive protein (CRP), interleukin (IL)-6, fibrinogen, blood viscosity, insulin, glycated haemoglobin (Hb<sub>A1c</sub>), and systolic (SBP) and diastolic blood pressure (DBP). PEL82 did not have data on metabolic measures from NMR platform and contributed data to analyses of conventional lipid risk factors (total cholesterol, HDL-c, LDL-c, and triglycerides (TG)), and some of the additional measures described (CRP, Hb<sub>A1c</sub>, SBP, DBP). Adiponectin was assayed using an enzyme-linked immunosorbent assay (ELISA) in PEL82, BWHHS and WHII. Data on adiponectin level was not available from CaPS, UKCTOCS, ALSPAC-M, and the Metabolomics consortium. Blood samples used for adiponectin, NMR metabolites, and other blood based outcomes were taken after overnight or minimum 6-hours fast in BWHHS, CaPS, and ALSPAC-M and on non-fasting samples in PEL82 and UKCTOCS. In WHII, participants attending the morning clinic were asked to fast overnight and those attending in the afternoon were asked to have a light, fat-free breakfast before 0800 hours. The vast majority of samples contributing to the Metabolomics consortium were fasting samples.

### ***Genotyping***

BWHHS, CaPS, WHII and UKCTOCS participants were genotyped using MetaboChip, a platform comprising 200,000 SNPs, which cover the loci identified by GWAS in cardio-metabolic diseases, and rare variants from the 1000 Genomes Project (36). Quality control criteria and imputation using 1000 Genomes European ancestry reference samples have been previously described for studies within UCLEB consortium (37). In ALSPAC-M, 557,124 SNPs were directly genotyped using Illumina human660W quad. For quality control, SNPs were excluded if missingness > 5%, Hardy-Weinberg equilibrium P-value <  $1 \times 10^{-6}$  or minor allele frequency < 1%, and samples were excluded if missingness > 5%, indeterminate X chromosome heterozygosity, extreme autosomal heterozygosity or showing evidence of population stratification. Imputation was performed using 1000 genomes reference panel (Phase 1, Version 3) (phased using Shapelt v2.r644, haplotype release date Dec 2013) and Impute V2.2.2. For PEL82, genotyping was performed by using the Illumina HumanOmni2.5-8v1 array (Illumina Inc.) and approximately 2,500,000 SNPs were genotyped (38). For PEL82, quality control criteria have been previously described (38) and imputation was performed in two steps: first, genotypes were phased using SHAPEIT; then, IMPUTE2 was used for the actual imputation. For autosomal and X-chromosome SNPs, 1000 Genomes Phase I integrated haplotypes (December 2013 release) and 1000 Genomes Phase I integrated variant set (March 2012 release), respectively, were used. For PEL82, ancestry-informative principal components were based on 370,539 SNPs shared by samples from the HapMap Project, the Human Genome Diversity Project (HGDP), and PEL82. The following HapMap samples were used as external panels: 266 Africans, 262 Europeans (American and Italian), 77 admixed Mexican Americans, 83 African Americans, and 93 Native Americans from the HGDP [more details can be found in (39)]. Cohorts contributing to the Metabolomics consortium used different SNP arrays, non-genotyped SNPs were imputed using a 1000 Genomes Project March 2012 version and SNPs with accurate imputation (proper info > 0.4) and minor allele count >3 were combined in fixed-effects meta-analysis using double genomic control correction. Further details can be found in the consortium publication (30).

### ***Other covariates***

Anthropometric variables (weight and height) were measured in each study using standard procedures and body mass index (BMI) was calculated as weight



(kg)/height (m)<sup>2</sup>. Demographic and smoking status information were obtained through questionnaires.

### **Data analysis**

Prior to analyses, metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. Pregnant women from PEL82 (n = 73) and ALSPAC-M (n = 12) were excluded. As the 74 analysed metabolites are highly correlated, we adopted a similar strategy to the Metabolomics consortium (30) to correct for multiple testing by estimating the number of independent tests as the number of principal components that explained over 95% of variance in metabolites concentration using data from the two studies (BWHHS and WHII) with the largest available number of metabolites (n = 27 principal components in both studies). As a result, for both multivariable and Mendelian randomization analyses, we corrected for multiple testing using the Bonferroni method considering 27 independent tests ( $p = 0.05 \div 27 \approx 0.0019$ ).

### Multivariable regression analysis

The conventional multivariable regression association of adiponectin with individual metabolites was estimated using a two-stage individual participant meta-analysis. In the first stage, linear regression models were fitted for each study. In the second stage, study-specific estimates were meta-analysed using DerSimonian & Laird random effect model (40). Heterogeneity across studies was assessed using  $I^2$  (as a measure of the relative size of between-study variation and within-study error) (41). Three types of subgroup analyses were conducted: sex-stratified analysis, analysis excluding individuals with high risk of cardiometabolic disease (those that had experienced coronary artery disease or stroke or those older than 65 years old) and analysis restricted to European studies (excluding PEL82).

### Genetic analyses

*Selection of genetic variants.* The SNPs used for the Mendelian randomization analysis were selected from 145 SNPs with good evidence ( $p < 5 \times 10^{-8}$ ) for

association with blood adiponectin concentration in the European ancestry GWAS meta-analysis from the ADIPOGen consortium (42). Independent SNPs within the *ADIPOQ* locus ( $\pm 50$  kb) have been previously selected by Dastani et al (2013) (43) by linkage disequilibrium (LD) pruning of the genome-wide significant SNPs, retaining SNPs that explained most variance in adiponectin concentration in each LD block (LD threshold:  $R^2 < 0.05$  in HapMap CEU population (Utah residents with Northern and Western European ancestry)). This resulted in four SNPs (rs6810075, rs16861209, rs17366568, and rs3774261), which are estimated to explain approximately 4% of variance in adiponectin concentration (**Table 2 and Supplementary methods**). Data for the association of each selected SNP with adiponectin concentration in the discovery GWAS sample was downloaded from ADIPOGen website (<https://www.mcgill.ca/genepi/adipogen-consortium>).

*Association of genetic variants with classical confounders.* The association between genetic variants and classical confounders [sex, age, ancestry (European vs non European), current smoking (yes vs no), and body mass index] was examined for each study using logistic or linear regression models for binary or continuous variables, respectively.

*Mendelian randomization analysis.* In order to allow all participants with relevant genetic and metabolic measure data to contribute to analyses, even when adiponectin data was not available (as in CaPS, UKCTOCS, ALSPAC-M, and Metabolomics consortium), a two-sample Mendelian randomization design was used, in which data for the association between genetic variants and adiponectin levels were obtained from an external data source, the ADIPOGen consortium (42). The two-sample Mendelian randomization is a recent extension to the more conventional one-sample Mendelian randomization and has the additional advantage of avoiding bias due to genetic variants correlating with confounders by chance (statistical overfitting) when samples are independent (44). The two-sample Mendelian randomization estimates and respective standard errors were obtained using the inverse variance-weighted (IVW) method, as described by Burgess et al. (45) and detailed in Supplementary Methods. Study-specific Mendelian randomization estimates were meta-analysed using DerSimonian & Laird random effect model (40). Heterogeneity across studies was assessed using  $I^2$  (41). Subgroup analyses were conducted considering individual-level (sex and risk of cardiometabolic disease) and study-level characteristics (European vs non-European studies). The Metabolomics

consortium did not contribute to subgroup analysis of individual-level characteristics as only summary data was available. Results from conventional multivariable and Mendelian randomization analyses were compared by using the Z-test for each metabolic measure (details in the Supplementary methods) and by estimating the correlation between multivariable and Mendelian randomization estimates across all metabolic measures. Power calculations for Mendelian randomization analysis are available in **Supplementary table 1**.

## RESULTS

The study included a median sample size of 3,006 adults in the multivariable analysis (range: 2,497-5,906) and a median sample size of 23,884 adults in the Mendelian randomization analysis (range: 4,645-38,058). Characteristics of participants from each contributing study are listed in **Table 2**.

### Adiponectin and the Systemic Metabolic Profile

In the multivariable analysis, adiponectin was associated with 59 out of 74 (80%) metabolites at nominal level ( $p < 0.05$ ) and 49 out of 74 (66%) after correcting for multiple testing ( $p < 0.0019$ ). Overall, higher circulating adiponectin was associated with a healthier systemic metabolite profile. Blood adiponectin concentration was strongly related to multiple lipoprotein traits. With higher adiponectin concentration, lipid concentration was lower in VLDL subclasses and higher in HDL subclasses, except for small HDL. There was no strong evidence of circulating adiponectin associating with total lipid content in LDL subclasses or in IDL, although adiponectin concentration was inversely associated with LDL-cholesterol. Higher adiponectin was associated with lower concentration of cholesterol and triglycerides, lower mean particle diameter in VLDL and higher cholesterol concentration and mean particle diameter in HDL. Higher adiponectin concentration was also associated with higher concentration of apolipoprotein (Apo)-AI and phospholipids and lower concentration of triglycerides and diglycerides (**Figure 2**).

Higher circulating adiponectin was also associated with healthier glycemic status (lower glucose and insulin concentration), lower blood concentration of glycolysis-related metabolites (lactate and pyruvate), saturated fatty acids, systemic inflammatory markers (C-reactive protein, fibrinogen, interleukin-6, glycoprotein acetyls and blood viscosity), systolic blood pressure, creatinine, and higher ketone bodies (acetoacetate). In addition, higher adiponectin concentration was associated with lower concentrations of free branched chain amino acids (isoleucine, leucine, and valine), aromatic amino acids (phenylalanine and tyrosine), and alanine and higher concentration of glutamine (**Figure 3**).

In the multivariable analyses, evidence of heterogeneity in pooled estimates across studies was substantial ( $I^2 = 50\%-75\%$ ) for 12 and very high ( $I^2 > 75\%$ ) for 15 metabolic measures (**Figure 2** and **3** and **Supplementary table 2**). This did not

seem to be accounted by sex (**Supplementary figures 1 to 4**), geographic location (**Supplementary figures 5 and 6**), or high risk of disease (**Supplementary figures 7 and 8**).

### **Causal effects of adiponectin on the Systemic Metabolic Profile**

Characteristics of the four SNPs (rs6810075, rs16861209, rs17366568 and rs3774261) used in Mendelian randomization and their association with adiponectin concentration are shown in **Table 3**. Overall, SNPs effect allele frequency was similar across studies. Two SNPs had lower allele frequency in the Metabolomics consortium (rs6810075: 51% vs. 65-69% in other studies; rs16861209: 5% vs. 9-11% in other studies) and one SNP had a higher frequency in PEL82 compared to other studies (rs3774261: 49% vs. 38-39% in other studies) (**Table 3**). As expected, the selected SNPs were not associated with classical confounders overall (**Supplementary table 3**).

Findings from Mendelian randomization analysis were largely inconsistent with results from multivariable analysis. Firstly, there was no evidence that adiponectin influenced HDL and VLDL traits (**Figure 2**). Secondly, genetically-increased adiponectin levels were not associated with glycemic traits, free amino acids, and glycolysis-related metabolites (**Figure 3**). Results were less conclusive for some inflammatory markers (IL-6 and fibrinogen) (**Figure 3**). Thirdly, there was strong statistical evidence that associations from multivariable and Mendelian randomization analyses were inconsistent with each other (**Figure 2** and **Figure 3**) and the overall correlation between multivariable and Mendelian randomization estimates was very low ( $r = 0.10$ ) (**Figure 4**). Finally, in the Mendelian randomization analysis, adiponectin was not associated with any of the metabolic analyses at either  $p < 0.05$  or  $p < 0.00068$ .

In the Mendelian randomization analyses, evidence of heterogeneity in pooled estimates across studies were substantial ( $I^2 = 50\%-75\%$ ) for 14 and very high ( $I^2 > 75\%$ ) for 3 metabolic measures, suggesting lower heterogeneity in models from genetic analysis than from the multivariable analyses (**Figure 2** and **3** and **Supplementary table 2**). This did not seem to be driven by sex differences (**Supplementary figures 1 to 4**), geographic location/ethnicity (**Supplementary figures 5 and 6**), or high risk of disease (**Supplementary figures 7 and 8**).

## DISCUSSION

In up to 5,906 adults we found using multivariable regression analyses that circulating adiponectin was associated with a pattern of systemic metabolites levels associated with good health. Higher blood adiponectin concentration was associated with higher HDL lipids and lower VLDL lipids, glycaemia, and branched-chain amino acids levels. However, when we used genetic variants in the *ADIPOQ* locus to test the causal effect of adiponectin on systemic metabolic profiles amongst up to 38,058 adults, we found little evidence that the associations were causal.

Despite the evidence of shared genetic architecture between adiponectin concentration and cardio-metabolic diseases (42), previous Mendelian randomization studies have cast doubt on the causal role of blood adiponectin levels in the risk of type 2 diabetes (18) and coronary heart disease (21). In addition, there seems to be no consistent evidence that circulating adiponectin causally affects traditional cardiovascular risk factors, such as HDL-c, LDL-c, triglycerides, and fasting glucose in the population (18). We have added importantly to those previous studies and explored effects on systemic metabolic profiles. Taken together, this and previous Mendelian randomization studies suggest that the association between circulating adiponectin and metabolic biomarkers and cardio-metabolic diseases is likely to be explained by shared factors (confounding) rather than by a direct role of adiponectin on metabolism and downstream cardio-metabolic disease. These results are in contrast to findings from animal models pointing to insulin-sensitizing, and anti-atherogenic actions of adiponectin (1).

Circulating adiponectin is known to be substantially reduced among obese individuals, particularly in the presence of central fat accumulation (46). A recent Mendelian randomization study examining the causal metabolic effects of BMI demonstrated that lower BMI was related to favorable lipoprotein subclass profile and lower concentration of branched-chain amino acids, inflammatory markers, and insulin (35), which is remarkably similar to our results from the conventional multivariable analysis. In addition, numerous studies have shown that adiponectin production is suppressed by insulin action in humans, which seems to be at least partly attributed to regulation at the transcriptional level (11, 47). As an example, elevated circulating adiponectin is found in contexts of both primary deficiency of insulin (type 1 diabetes) (48) and global insulin resistance due to genetic or acquired defects in the insulin receptor (49). Evidence from animal models has raised the

possibility of a bidirectional relationship between adiponectin and insulin concentration (50). Early Mendelian randomization studies did indicate that adiponectin could mitigate insulin resistance (19, 20); however, these results could not be replicated in a larger Mendelian randomization study (18), as well as in our study presented here. The well-known metabolic effects of adiposity and insulin on circulating adiponectin concentration reinforce that the clustering of adiponectin and several traditional and novel biomarkers is likely to result from confounding due to increasing adiposity and disruption of insulin action.

Strengths of our study include detailed metabolic profile in several longitudinal studies, which enabled us to characterize the metabolic profile of high adiponectin concentration beyond traditional biomarkers, as well as the use of Mendelian randomization to disentangle the causal effect of adiponectin on the metabolism. Mendelian randomization analysis can reliably test for the presence of a causal relation under the three assumptions of an instrumental variable that the genetic variants are robustly associated with the risk factor of interest (adiponectin) (1), should only affect the outcome (metabolites) through the exposure (2), and are not associated with exposure-outcome confounders (3) (51). To ensure that IV assumptions were met, or were at least plausible, we only used SNPs strongly and specifically (within *ADIPOQ* gene) related to adiponectin concentration as instrumental variables and we adjusted for population structure in models using data from PEL82 to avoid confounding by population stratification. One of the limitations of our study was the limited power in subgroup analyses including only individual-level data (sex- and risk-stratified analyses), which limited our investigation of potential sources of heterogeneity. Another limitation was the absence of data on high-molecular weight adiponectin, which is believed to account for most of the adiponectin biological effects in experimental settings. However, most human (and many animal model) studies have not used high-molecular weight adiponectin, and we found the same multivariable observational associations as in previous studies.

Overall, our findings suggest that altered total blood adiponectin concentration is an epiphenomenon in the context of metabolic disease, rather than a key determinant. Therefore, interventions targeting manipulation of adiponectin concentration are unlikely to result in therapeutic benefits for tackling cardiovascular diseases. Our results highlight the potential of Mendelian randomization analysis and high-throughput metabolomics profiling to yield important insights to advance our



understanding in the pathophysiology of common complex diseases and to inform which targets are 'best-bets' for taking forward into drug development, given that drug target validation is a key obstacle underlying the unsustainably high rate of drug development failure. Whilst our, and other studies, suggest adiponectin is not a valuable target for developing drugs aimed at preventing cardio-metabolic diseases, it may nonetheless be a valuable biomarker for predicting these diseases given the wide ranging associations shown here. The associations we have found would need to be replicated in additional independent studies before testing their ability to predict disease outcomes.

## Acknowledgements

We acknowledge Andy Ryan for his contribution to data collection from UKCTOCS. MCB, DLSF, DAL and TRG work in the MRC Integrative Epidemiology Unit at the University of Bristol that receives funding from the UK Medical Research Council (MC\_UU\_12013/5 and MC\_UU\_12013/8). DAL is a UK National Institute of Health Research Senior Investigator (NF-SI-0611-10196). MKiv is supported by the UK Medical Research Council (K013351). **PEL82** (1982 Pelotas Birth Cohort) is conducted by Postgraduate Program in Epidemiology at Universidade Federal de Pelotas with the collaboration of the Brazilian Public Health Association (ABRASCO). From 2004 to 2013, the Wellcome Trust supported PEL82. The International Development Research Center, World Health Organization, Overseas Development Administration, European Union, National Support Program for Centers of Excellence (PRONEX), the Brazilian National Research Council (CNPq), and the Brazilian Ministry of Health supported previous phases of the study. The **UCLEB** (UCL-LSHTM-Edinburgh-Bristol) consortium, which is supported by BHF Programme Grant RG/10/12/28456, consists of 12 studies: Northwick Park Heart Study II (NPHS II), British Regional Heart Study (BRHS), Whitehall II Study (WHII), English Longitudinal Study of Ageing (ELSA), Medical Research Council National Survey of Health and Development (MRC NSHD), 1958 Birth cohort (1958BC), Caerphilly prospective study (CaPS), British Women's Heart and Health Study (BWHHS), Edinburgh Artery Study (EAS), Edinburgh Heart Disease Prevention Study (EHDPS), Edinburgh Type 2 Diabetes Study (ET2DS) and Asymptomatic Atherosclerosis Aspirin Trial (AAAT).



**BWHHS** is supported by funding from the British Heart Foundation and the Department of Health Policy Research Programme (England). EAS is funded by the British Heart Foundation (Programme Grant RG/98002), with MetaboChip genotyping funded by a project grant from the Chief Scientist Office of Scotland (Project Grant CZB/4/672). The **WHII study** is supported by grants from the Medical Research Council (K013351), British Heart Foundation (RG/07/008/23674), Stroke Association, the US National Heart Lung and Blood Institute (5RO1 HL036310), the US National Institute on Aging (5RO1AG13196) the US Agency for Healthcare Research and Quality (HS06516); and the John D. and Catherine T. MacArthur Foundation Research Networks on Successful Midlife Development and Socio-economic Status and Health. **CaPS** was funded by the Medical Research Council and undertaken by the former MRC Epidemiology Unit (South Wales). The CaPS DNA bank was established with funding from a MRC project grant. The CaPS data archive is maintained by the University of Bristol. MRC Integrative Epidemiology Unit, Bristol is supported by MRC grants (MR\_UU\_12013/1, MR\_UU\_12013/5 and MR\_UU\_12013/8). **UKCTOCS** was funded by the Medical Research Council (G9901012 and G0801228), Cancer Research UK (C1479/A2884), and the Department of Health, with additional support from The Eve Appeal. Phenotypic data for this case control dataset was supported by the National Institute for Health Research, Biomedical Research Centre at University College London Hospital. **ALSPAC-M** phenotypic data was collected with funding from the British Heart Foundation (SP/07/008/24066), Wellcome Trust (WT092830M) and UK Research Councils (UKRC) via the MRC (G1001357); genetic data collection was funded by the Wellcome Trust (WT088806). In addition the ALSPAC full study receives core support from The University of Bristol, UK Medical Research Council and the Wellcome Trust (102215/2/13/2) and the University of Bristol. The ALSPAC team is extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

Summary genome-wide association data on adiponectin have been contributed by ADIPOGen Consortium and have been downloaded from <https://www.mcgill.ca/genepi/adipogen-consortium>. Summary genome-wide

association data on metabolic measures have been contributed by Kettunen et al. (30) and have been downloaded from <https://www.mcgill.ca/genepi/adipogen-consortium>.

### **Conflicts of interest**


No competing interests: MCB, AJDB, DLSF, JPC, BLH, MKiv, MKu, TRG, YBS, DFF, IOO, AGM, EF, DAL, and ADH. UM has stock ownership in and research funding from Abcodia Pvt Ltd.

## REFERENCES

1. Turer AT, Scherer PE. Adiponectin: mechanistic insights and clinical implications. *Diabetologia*. 2012;55(9):2319-26.
2. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*. 1999;257(1):79-83.
3. Li S, Shin HJ, Ding EL, van Dam RM. Adiponectin Levels and Risk of Type 2 Diabetes A Systematic Review and Meta-analysis. *Jama-Journal of the American Medical Association*. 2009;302(2):179-88.
4. Polyzos SA, Toulis KA, Goulis DG, Zavos C, Kountouras J. Serum total adiponectin in nonalcoholic fatty liver disease: a systematic review and meta-analysis. *Metabolism*. 2011;60(3):313-26.
5. Matsuzawa Y, Funahashi T, Kihara S, Shimomura I. Adiponectin and metabolic syndrome. *Arterioscler Thromb Vasc Biol*. 2004;24(1):29-33.
6. Wannamethee SG, Welsh P, Whincup PH, Sawar N, Thomas MC, Gudnarsson V, et al. High adiponectin and increased risk of cardiovascular disease and mortality in asymptomatic older men: does NT-proBNP help to explain this association? *Eur J Cardiovasc Prev Rehabil*. 2011;18(1):65-71.
7. Lee ES, Park S-s, Kim E, Yoon YS, Ahn H-Y, Park C-Y, et al. Association between adiponectin levels and coronary heart disease and mortality: a systematic review and meta-analysis. *International Journal of Epidemiology*. 2013;42(4):1029-39.
8. Menon V, Li L, Wang X, Greene T, Balakrishnan V, Madero M, et al. Adiponectin and mortality in patients with chronic kidney disease. *J Am Soc Nephrol*. 2006;17(9):2599-606.
9. Beatty AL, Zhang MH, Ku IA, Na B, Schiller NB, Whooley MA. Adiponectin is associated with increased mortality and heart failure in patients with stable ischemic heart disease: data from the Heart and Soul Study. *Atherosclerosis*. 2012;220(2):587-92.
10. Sattar N, Nelson SM. Adiponectin, diabetes, and coronary heart disease in older persons: unraveling the paradox. *J Clin Endocrinol Metab*. 2008;93(9):3299-301.
11. Cook JR, Semple RK. Hypoadiponectinemia--cause or consequence of human "insulin resistance"? *J Clin Endocrinol Metab*. 2010;95(4):1544-54.
12. Lawlor DA, Harbord RM, Sterne JA, Timpson N, Davey Smith G. Mendelian randomization: using genes as instruments for making causal inferences in epidemiology. *Stat Med*. 2008;27(8):1133-63.
13. Smith GD, Lawlor DA, Harbord R, Timpson N, Day I, Ebrahim S. Clustered environments and randomized genes: a fundamental distinction between conventional and genetic epidemiology. *PLoS Med*. 2007;4(12):e352.
14. Ference BA, Julius S, Mahajan N, Levy PD, Williams KA, Flack JM. Clinical effect of naturally random allocation to lower systolic blood pressure beginning before the development of hypertension. *Hypertension*. 2014;63(6):1182-8.
15. Collaboration CRPCHDG, Wensley F, Gao P, Burgess S, Kaptoge S, Di Angelantonio E, et al. Association between C reactive protein and coronary heart disease: mendelian randomisation analysis based on individual participant data. *BMJ*. 2011;342:d548.
16. Collaboration IRGCERF, Sarwar N, Butterworth AS, Freitag DF, Gregson J, Willeit P, et al. Interleukin-6 receptor pathways in coronary heart disease: a collaborative meta-analysis of 82 studies. *Lancet*. 2012;379(9822):1205-13.
17. Swerdlow DI, Preiss D, Kuchenbaecker KB, Holmes MV, Engmann JE, Shah T, et al. HMG-coenzyme A reductase inhibition, type 2 diabetes, and bodyweight: evidence from genetic analysis and randomised trials. *Lancet*. 2015;385(9965):351-61.
18. Yaghoobkar H, Lamina C, Scott RA, Dastani Z, Hivert MF, Warren LL, et al. Mendelian randomization studies do not support a causal role for reduced circulating adiponectin levels in insulin resistance and type 2 diabetes. *Diabetes*. 2013;62(10):3589-98.

19. Mente A, Meyre D, Lanktree MB, Heydarpour M, Davis AD, Miller R, et al. Causal relationship between adiponectin and metabolic traits: a Mendelian randomization study in a multiethnic population. *PLoS One*. 2013;8(6):e66808.
20. Gao H, Fall T, van Dam RM, Flyvbjerg A, Zethelius B, Ingelsson E, et al. Evidence of a Causal Relationship Between Adiponectin Levels and Insulin Sensitivity. *Diabetes*. 2013;62(4):1338-44.
21. Borges MC, Lawlor DA, de Oliveira C, White J, Horta BL, Barros AJ. Role of Adiponectin in Coronary Heart Disease Risk: A Mendelian Randomization Study. *Circ Res*. 2016;119(3):491-9.
22. Würtz P, Mäkinen VP, Soininen P, Kangas AJ, Tukiainen T, Kettunen J, et al. Metabolic signatures of insulin resistance in 7,098 young adults. *Diabetes*. 2012;61(6):1372-80.
23. Horta BL, Gigante DP, Gonçalves H, dos Santos Motta J, Loret de Mola C, Oliveira IO, et al. Cohort Profile Update: The 1982 Pelotas (Brazil) Birth Cohort Study. *Int J Epidemiol*. 2015;44(2):441, a-e.
24. Victora CG, Barros FC. Cohort profile: The 1982 Pelotas (Brazil) Birth Cohort Study. *International Journal of Epidemiology*. 2006;35(2):237-42.
25. Lawlor DA, Ebrahim S, Davey Smith G, study Bwshah. Socioeconomic position in childhood and adulthood and insulin resistance: cross sectional survey using data from British women's heart and health study. *BMJ*. 2002;325(7368):805.
26. Marmot M, Brunner E. Cohort Profile: the Whitehall II study. *Int J Epidemiol*. 2005;34(2):251-6.
27. Patterson CC, Blankenberg S, Ben-Shlomo Y, Heslop L, Bayer A, Lowe G, et al. Which biomarkers are predictive specifically for cardiovascular or for non-cardiovascular mortality in men? Evidence from the Caerphilly Prospective Study (CaPS). *Int J Cardiol*. 2015;201:113-8.
28. Menon U, Gentry-Maharaj A, Ryan A, Sharma A, Burnell M, Hallett R, et al. Recruitment to multicentre trials--lessons from UKCTOCS: descriptive study. *BMJ*. 2008;337:a2079.
29. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al. Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol*. 2013;42(1):97-110.
30. Kettunen J, Demirkan A, Wurtz P, Draisma HH, Haller T, Rawal R, et al. Genome-wide study for circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. *Nat Commun*. 2016;7:11122.
31. Würtz P, Havulinna AS, Soininen P, Tynkkynen T, Prieto-Merino D, Tillin T, et al. Metabolite profiling and cardiovascular event risk: a prospective study of 3 population-based cohorts. *Circulation*. 2015;131(9):774-85.
32. Würtz P, Wang Q, Soininen P, Kangas AJ, Fatemifar G, Tynkkynen T, et al. Metabolomic Profiling of Statin Use and Genetic Inhibition of HMG-CoA Reductase. *J Am Coll Cardiol*. 2016;67(10):1200-10.
33. Soininen P, Kangas AJ, Wurtz P, Tukiainen T, Tynkkynen T, Laatikainen R, et al. High-throughput serum NMR metabolomics for cost-effective holistic studies on systemic metabolism. *Analyst*. 2009;134(9):1781-5.
34. Soininen P, Kangas AJ, Wurtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ Cardiovasc Genet*. 2015;8(1):192-206.
35. Wurtz P, Wang Q, Kangas AJ, Richmond RC, Skarp J, Tiainen M, et al. Metabolic signatures of adiposity in young adults: Mendelian randomization analysis and effects of weight change. *PLoS Med*. 2014;11(12):e1001765.
36. Voight BF, Kang HM, Ding J, Palmer CD, Sidore C, Chines PS, et al. The metabochip, a custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric traits. *PLoS Genet*. 2012;8(8):e1002793.
37. Shah T, Engmann J, Dale C, Shah S, White J, Giambartolomei C, et al. Population genomics of cardiometabolic traits: design of the University College London-London School of Hygiene and Tropical Medicine-Edinburgh-Bristol (UCLEB) Consortium. *PLoS One*. 2013;8(8):e71345.

38. Borges MC, Hartwig FP, Oliveira IO, Horta BL. Is there a causal role for homocysteine concentration in blood pressure? A Mendelian randomization study. *Am J Clin Nutr.* 2016;103(1):39-49.
39. Lima-Costa MF, Rodrigues LC, Barreto ML, Gouveia M, Horta BL, Mambrini J, et al. Genomic ancestry and ethnoracial self-classification based on 5,871 community-dwelling Brazilians (The Epigen Initiative). *Sci Rep.* 2015;5:9812.
40. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7(3):177-88.
41. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21(11):1539-58.
42. Dastani Z, Hivert MF, Timpson N, Perry JR, Yuan X, Scott RA, et al. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet.* 2012;8(3):e1002607.
43. Dastani Z, Johnson T, Kronenberg F, Nelson CP, Assimes TL, März W, et al. The shared allelic architecture of adiponectin levels and coronary artery disease. *Atherosclerosis.* 2013;229(1):145-8.
44. Lawlor DA. Commentary: On Gao C et al. Mendelian randomization study of adiposity-related traits and risk of breast, ovarian, prostate, lung and colorectal cancer. *Int J Epidemiol.* 2016.
45. Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol.* 2013;37(7):658-65.
46. Borges M, Oliveira I, Freitas D, Horta B, Ong K, Gigante D, et al. Obesity-induced hypoadiponectinemia: the opposite influence of central and peripheral fat compartments. *International Journal of Epidemiology* (under review). 2016.
47. Kern PA, Di Gregorio GB, Lu T, Rassouli N, Ranganathan G. Adiponectin expression from human adipose tissue: relation to obesity, insulin resistance, and tumor necrosis factor- $\alpha$  expression. *Diabetes.* 2003;52(7):1779-85.
48. Imagawa A, Funahashi T, Nakamura T, Moriwaki M, Tanaka S, Nishizawa H, et al. Elevated serum concentration of adipose-derived factor, adiponectin, in patients with type 1 diabetes. *Diabetes Care.* 2002;25(9):1665-6.
49. Semple RK, Cochran EK, Soos MA, Burling KA, Savage DB, Gorden P, et al. Plasma adiponectin as a marker of insulin receptor dysfunction: clinical utility in severe insulin resistance. *Diabetes Care.* 2008;31(5):977-9.
50. Kubota N, Terauchi Y, Kubota T, Kumagai H, Itoh S, Satoh H, et al. Pioglitazone ameliorates insulin resistance and diabetes by both adiponectin-dependent and -independent pathways. *J Biol Chem.* 2006;281(13):8748-55.
51. Swerdlow DI, Kuchenbaecker KB, Shah S, Sofat R, Holmes MV, White J, et al. Selecting instruments for Mendelian randomization in the wake of genome-wide association studies. *Int J Epidemiol.* 2016.
52. Shim H, Chasman DI, Smith JD, Mora S, Ridker PM, Nickerson DA, et al. A multivariate genome-wide association analysis of 10 LDL subfractions, and their response to statin treatment, in 1868 Caucasians. *PLoS One.* 2015;10(4):e0120758.



**Figure 1.** Schematic representation of studies contributing to each analytical approach.

From the available data sources, three had data on adiponectin and metabolic measures and could contribute to multivariable analysis (PEL82, BWHHS and WHII) and all had data on genetic variants and metabolic measures and could contribute to Mendelian randomization analysis (PEL82, BWHHS, WHII, CaPS, UKCTOCS, ALSPAC-M, and Metabolomics consortium). ALSPAC-M: The Avon Longitudinal Study of Children and Parents – mothers’ cohort; BWHHS: British Women’s Heart and Health Study; CaPS: The Caerphilly Prospective Study; PEL82: 1982 Pelotas Birth Cohort; UKCTOCS: case-control study nested in The United Kingdom Collaborative Trial of Ovarian Cancer Screening; WHII: Whitehall-II Study.

**Table 1. Characteristics of participating studies**

	PEL82	BWHHS	WHII	CaPS	UKCTOCS case-control <sup>a</sup>	ALSPAC-M	Metabolomics consortium
Study design	Cohort	Cohort	Cohort	Cohort	Nested case-control study	Cohort	14 cohorts
Setting	Brazil	United Kingdom	United Kingdom	United Kingdom	United Kingdom	United Kingdom	Europe
Recruitment setting	Hospitals	General practices	Workplace	General practices and electoral register	Hospitals	Media information, community locations, and health services	Multiple settings
Participants	Adults aged 30 years old born in the city of Pelotas in 1982	Women aged 60-79 years old at recruitment	Civil servants aged 45-69 years at phase 5	Men aged 52-72 years old at phase III	Postmenopausal women aged 50 years old and above at recruitment	Women aged 34-63 years old residing in a defined area in the South West of England that gave birth between 01/04/1991 to 31/12/1992	Adults recruited for multiple studies (mean age: 45 years old)
Phase of data collection	2012 follow-up	Recruitment (1999-2001)	Phase 5 (1997-1999)	Phase III (1989-1993)	Recruitment (2001-2005)	Follow-up clinic assessment (2009-2011)	Data collected in different phases according to each study
Blood samples fasted	No	Yes	Mixed	Yes	No	Yes	Yes (for the vast majority of blood samples)
N at data collection phase	3,701	4,286	7,870	2,154	4,867	4,834	25,072
N with adiponectin	3,541	498	2,662	0	0	0	0
N with metabolites	3,530 <sup>b</sup>	3,780	4,641	1,225	4,813	4,138	25,072
N with other phenotypes <sup>c</sup>	3,530-3,617	3,636-3,964	4,620-4,874	608-1,207	0	4,092-4,568	—
N with genotype	2,898	1,980	3,078	1,349	1,472	8,672 <sup>d</sup>	25,072
N for multivariable associations	2,753-2,762	396-497	2,442-2,656	—	—	—	—
N for MR analyses	2,753-2,783	1,656-1,967	2,773-3,020	101-1,211	1,067-1,435	2,548-3,375	12,978-24,924
Website	<a href="http://www.epidemiologyofpelotas.org.br/site/content/portal_1982-en/index.php">http://www.epidemiologyofpelotas.org.br/site/content/portal_1982-en/index.php</a>	<a href="http://www.lshtm.ac.uk/epidemiology/research/bwhhs/index.html">http://www.lshtm.ac.uk/epidemiology/research/bwhhs/index.html</a>	<a href="http://www.ucl.ac.uk/whitehall">http://www.ucl.ac.uk/whitehall</a>	<a href="http://www.bristol.ac.uk/social-community-medicine/projects/caerphilly/about/">http://www.bristol.ac.uk/social-community-medicine/projects/caerphilly/about/</a>	<a href="http://www.instituteforwomenshealth.ucl.ac.uk/womens-cancer/gcra/ukctocs">http://www.instituteforwomenshealth.ucl.ac.uk/womens-cancer/gcra/ukctocs</a>	<a href="http://www.bristol.ac.uk/alspac/">http://www.bristol.ac.uk/alspac/</a>	<a href="http://www.computationalmedicine.fi/data/NMR_GWAS/">http://www.computationalmedicine.fi/data/NMR_GWAS/</a>

<sup>a</sup> The nested case-control study consisted of a subsample (n = 4,867) of the original UKCTOCS randomized controlled trial (N = 202,638 recruited individuals)

<sup>b</sup> For PEL82, the only metabolites available were glucose, total cholesterol, HDL-c, LDL-c and TAG

<sup>c</sup> Other phenotypes include systolic and diastolic blood pressure, glycated hemoglobin, C reactive protein, interleukin-6, fibrinogen and blood viscosity

<sup>d</sup> DNA samples were collected for the whole cohort in prior phases of ALSPAC-M cohort

ALSPAC-M: The Avon Longitudinal Study of Children and Parents – mothers' cohort; BWHHS: British Women's Heart and Health Study; CaPS: The Caerphilly Prospective Study; PEL82: 1982 Pelotas Birth Cohort; UKCTOCS: case-control study nested in The United Kingdom Collaborative Trial of Ovarian Cancer Screening; WHII: Whitehall-II Study. MR: Mendelian randomization analysis.

**Table 2.** Characteristics of studies' populations

	PEL82	BWHHS	WHII	CaPS	UKCTOCS	ALSPAC-M	Metabolomics consortium
	%						
Male	49	0	72	100	0	0	45
White	75	100	93	100	97	97	NA <sup>a</sup>
Smoker	24	12	17	20	—	11	NA
Overweight/obese	58	72	57	69	60	56	NA
	Median (p25, p75)						
Age (years)	30 (30, 30)	69 (64, 73)	55 (51, 61)	56 (53, 60)	66 (60, 70)	48 (45, 51)	45 (24, 61) <sup>b</sup>
Adiponectin (µg/mL)	7.9 (5.2, 11.9)	15.8 (10.8, 21.5)	8.5 (6.1, 12)	—	—	—	—
Glucose (mmol/L)	4.8 (4.4, 5.3)	4.7 (4.3, 5.1)	5 (4.7, 5.4)	3.8 (3.5, 4.2)	2.2 (1.7, 3.1)	4.4 (4.1, 4.7)	NA
HDL-c (mmol/L)	1.5 (1.2, 1.7)	1.6 (1.4, 1.9)	1.5 (1.3, 1.7)	0.9 (0.7, 1)	1.6 (1.4, 1.9)	1.7 (1.5, 1.9)	NA
LDL-c (mmol/L)	2.7 (2.3, 3.3)	2.3 (1.9, 2.8)	1.9 (1.6, 2.2)	1.6 (1.3, 1.9)	1.8 (1.4, 2.2)	1.5 (1.2, 1.8)	NA
TG (mmol/L)	1.1 (0.8, 1.6)	1.5 (1.1, 2)	1.1 (0.9, 1.5)	1.5 (1.2, 2)	1.5 (1.1, 2.1)	0.9 (0.7, 1.2)	—
SBP (mmHg)	120 (112, 130)	146 (130, 163)	121 (111, 133)	144 (130, 160)	—	117 (110, 125)	—
DBP (mmHg)	75 (69, 81)	79 (71, 87)	77 (70, 84)	84 (76, 92)	—	71 (66, 77)	—

<sup>a</sup> Cohorts contributing to the Metabolomics consortium were of European origin

<sup>b</sup> Overall mean age (and range of mean age across studies)


ALSPAC-M: The Avon Longitudinal Study of Children and Parents – mothers' cohort; BWHHS: British Women's Heart and Health Study; CaPS: The Caerphilly Prospective Study; DBP: diastolic blood pressure; HDL-c: high-density lipoprotein-cholesterol; LDL-c: low-density lipoprotein-cholesterol; PEL82: 1982 Pelotas Birth Cohort; SBP: systolic blood pressure; TG: triglycerides; UKCTOCS: case-control study nested in The United Kingdom Collaborative Trial of Ovarian Cancer Screening; WHII: Whitehall-II Study.



**Table 3.** Characteristics of SNPs selected for Mendelian randomization analysis


	SNP			
	rs6810075	rs16861209	rs17366568	rs3774261
Chr	3	3	3	3
Position†	186548565	186563114	186570453	186571559
Closest gene	ADIPOQ	ADIPOQ	ADIPOQ-AS1, ADIPOQ	ADIPOQ-AS1, ADIPOQ
EA	T	A	G	A
NEA	C	C	A	G
<i>ADIPOGen consortium</i>				
EAF‡	0.63	0.07	0.90	0.39
Beta§	0.11	0.31	0.25	0.11
SE	0.01	0.02	0.01	0.01
<i>PEL82</i>				
EAF	0.65	0.11	0.92	0.49
Beta	0.13	0.33	0.22	0.08
SE	0.03	0.04	0.05	0.03
R <sup>2</sup>	0.008	0.021	0.005	0.002
<i>BWHHS</i>				
EAF	0.67	0.09	0.89	0.38
Beta	0.32	0.30	1.04	0.30
SE	0.10	0.14	0.24	0.08
R <sup>2</sup>	0.022	0.020	0.051	0.044
<i>WHII</i>				
EAF	0.68	0.10	0.89	0.38
Beta	0.16	0.36	0.56	0.14
SE	0.04	0.05	0.08	0.03
R <sup>2</sup>	0.008	0.027	0.025	0.010
<i>CaPS</i>				
EAF	0.69	0.10	0.89	0.39
<i>UKCTOCS</i>				
EAF	0.69	0.10	0.89	0.38
<i>ALSPAC-M</i>				
EAF	0.66	0.09	0.93	0.38
<i>Metabolomics consortium</i>				
EAF	0.51	0.05	0.88	0.36

Beta (and SE) refers to mean difference in standardized log adiponectin per additional SNP effect allele. †Genome Reference Consortium Human Build 37. ‡Extracted from Dastani et al (2012) (42). \*For CaPS, UKCTOCS and ALSPAC-M, data on adiponectin levels were not available. Chr: chromosome; EA: effect allele; NEA: non effect allele; EAF: effect allele frequency; SE: standard error; SNP: single nucleotide polymorphism.




**Figure 2.** Association of lipoprotein traits with blood adiponectin levels from observational and Mendelian randomization (MR) analysis.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, C: The Caerphilly Prospective Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, M: Metabolomics consortium, SD units: standard deviation units, CI: confidence interval.



**Figure 3.** Association of multiple metabolic measures with blood adiponectin levels from observational and Mendelian randomization analysis.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (\*\*\*\*) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, FAw6: omega-6 fatty acid, LA: linoleic acid, FAw3: omega-3 fatty acid, DHA: docosaexaenoic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, C: The Caerphilly Prospective Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, M: Metabolomics consortium, SD units: standard deviation units, CI: confidence interval.



**Figure 4.** Correlation between estimates from multivariable regression and Mendelian randomization (MR).

XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, FAW6: omega-6 fatty acid, LA: linoleic acid, DHA: docosaexaenoic acid, FAW3: omega-3 fatty acid, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, r: Pearson correlation coefficient.

## SUPPLEMENTARY METHODS

### Mendelian randomization analyses

The two-sample Mendelian randomization estimates and respective standard errors were obtained using the inverse variance-weighted (IVW) method with the following formulas:

$$\hat{\beta}_{IVW} = \frac{\sum_{k=1}^K X_k Y_k \sigma_{y_k}^{-2}}{\sum_{k=1}^K X_k^2 \sigma_{y_k}^{-2}} \quad SE_{\hat{\beta}_{IVW}} = \sqrt{\frac{1}{\sum_{k=1}^K X_k^2 \sigma_{y_k}^{-2}}}$$

Where  $X_k$  is the mean change in standardized log adiponectin units per additional effect allele of SNP  $k$  and  $Y_k$  is the mean change in standardized units of metabolic measures per additional effect allele of SNP  $k$  with standard error  $\sigma_{y_k}$ . To increase precision and avoid bias due to statistical overfitting, estimates for  $X_k$  were obtained from ADIPOGen consortium dataset (42). Prior to analysis, estimates from ADIPOGen consortium were standardized (converted from log adiponectin to standardized log adiponectin units) using individual level data from PEL82 with a similar adiponectin distribution (adiponectin concentration in ADIPOGen consortium: mean = 9.8  $\mu\text{g/ml}$  (standard deviation = 5.6); adiponectin concentration in 1982 Pelotas Birth Cohort: mean = 9.3  $\mu\text{g/ml}$  (standard deviation = 5.7)). Estimates for  $Y_k$  were derived from each study using linear regression models considering an additive model for SNP alleles.

### Comparison between multivariable and Mendelian randomization analyses

Results from conventional multivariable and Mendelian randomization analyses were compared using the Z-test:

$$Z = (\beta_{obs} - \beta_{MR}) / \sqrt{SE_{obs}^2 + SE_{MR}^2}$$

Where  $\beta_{obs}$  represents estimates from conventional observational analysis (with respective standard error,  $SE_{obs}$ ) and  $\beta_{MR}$  represents estimates from Mendelian randomization analysis (with respective standard error,  $SE_{MR}$ ).

## Proportion of variance in adiponectin concentration explained by genetic instruments

In order to estimate the strength of our genetic instruments, we estimated the phenotypic variance explained by a given SNP ( $R^2$ ) for adiponectin concentration. We used ADIPOGen summary data to approximate  $R^2$  for a given SNP based on the effect estimate for its association with the trait of interest (beta or  $\hat{\beta}$ ), respective standard error ( $se(\hat{\beta})$ ), minor allele frequency (MAF), and sample size (N). The following formula was used as previously described by Shim et al., 2015 (52):


$$R^2 \cong \frac{2\hat{\beta}^2 MAF(1 - MAF)}{2\hat{\beta}^2 MAF(1 - MAF) + (se(\hat{\beta}))^2 2NMAF(1 - MAF)}$$

The phenotypic variance explained by the composite genetic instrument (combining all SNPs) was estimated by the sum of SNP-specific  $R^2$  as shown below:

SNPs used as instrumental variables for adiponectin concentration in Mendelian randomization analysis and association with adiponectin concentration


rs ID	Chr	EA	NEA	EAF	$R^2$	N
rs6810075	3	T	C	0.63	0.0066	29140
rs16861209	3	A	C	0.01	0.0125	29199
rs17366568	3	G	A	0.91	0.0125	24865
rs3774261	3	A	G	0.60	0.0080	29081
Combined instrument	N/A	N/A	N/A	N/A	0.0396	N/A

Chr: chromosome; EA: effect allele (trait-increasing allele); NEA: non-effect allele;  $R^2$ : proportion of phenotypic variance explained by SNP; Beta: increase in standardized log adiponectin concentration per EA; SE: standard error; N: sample size; N/A: non applicable.



**Supplementary figure 1.** Association of lipoprotein traits with blood adiponectin levels from observational and Mendelian randomization (MR) analysis among women.


Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, , A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, SD units: standard deviation units, CI: confidence interval.



**Supplementary figure 2.** Association of multiple metabolic measures with blood adiponectin levels from observational and Mendelian randomization analysis among women.


Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks ("\*\*\*\*") if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  ("+") or very high if  $I^2 > 75\%$  ("++"). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column "MR vs MV (P-value)". Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, Faw6: omega-6 fatty acid, LA: linoleic acid, Faw3: omega-3 fatty acid, DHA: docosahexaenoic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers' cohort, SD units: standard deviation units, CI: confidence interval.






**Supplementary figure 3.** Association of lipoprotein traits with blood adiponectin levels from observational and Mendelian randomization (MR) analysis among men.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”), very high if  $I^2 > 75\%$  (“++”) or not applicable (“NA”) when only one study contributed to the estimate. P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MR (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, P: 1982 Pelotas Birth Cohort, W: Whitehall II Study, C: The Caerphilly Prospective Study, SD units: standard deviation units, CI: confidence interval.




**Supplementary figure 4.** Association of multiple metabolic measures with blood adiponectin levels from observational and Mendelian randomization analysis among men.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”) or not applicable (“NA”) when only one study contributed to the estimate. P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, Faw6: omega-6 fatty acid, LA: linoleic acid, Faw3: omega-3 fatty acid, DHA: docosaenoic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, P: 1982 Pelotas Birth Cohort, W: Whitehall II Study, C: The Caerphilly Prospective Study, SD units: standard deviation units, CI: confidence interval.




**Supplementary figure 5.** Association of lipoprotein traits with blood adiponectin levels from observational and Mendelian randomization (MR) analysis restricted to individuals of European ancestry.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, M: Metabolomics consortium, SD units: standard deviation units, CI: confidence interval.




**Supplementary figure 6.** Association of multiple metabolic measures with blood adiponectin levels from observational and Mendelian randomization analysis restricted to individuals of European ancestry.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (\*\*\*\*) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, FAw6: omega-6 fatty acid, LA: linoleic acid, FAw3: omega-3 fatty acid, DHA: docosaeic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, M: Metabolomics consortium, SD units: standard deviation units, CI: confidence interval.



**Supplementary figure 7.** Association of lipoprotein traits with blood adiponectin levels from observational and Mendelian randomization (MR) analysis among younger individuals (< 65 years old) free from cardiovascular disease.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, SD units: standard deviation units, CI: confidence interval.



**Supplementary figure 8.** Association of multiple metabolic measures with blood adiponectin levels from observational and Mendelian randomization analysis among younger individuals (< 65 years old) free from cardiovascular disease.

Values are expressed as units of standardized log metabolite concentration (and 95% CI) per 1 unit increment of standardized log adiponectin levels. P-values for the association between adiponectin and metabolites are indicated by three asterisks (“\*\*\*”) if lower than Bonferroni-adjusted threshold (P-value < 0.00068). Heterogeneity was considered substantial if  $I^2 = 50-75\%$  (“+”) or very high if  $I^2 > 75\%$  (“++”). P-values for the comparison between multivariable and Mendelian randomization estimates are displayed in the column “MR vs MV (P-value)”. Metabolic measures were adjusted for age, sex, and, if applicable, place of recruitment (BWHHS and UKCTOCS) or principal components of genomic ancestry (PEL82 and some studies contributing to Metabolomics consortium) and the resulting residuals were transformed to normal distribution by inverse rank-based normal transformation. TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, Faw6: omega-6 fatty acid, LA: linoleic acid, Faw3: omega-3 fatty acid, DHA: docosaenoic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure, P: 1982 Pelotas Birth Cohort, B: British Women Heart and Health Study, W: Whitehall II Study, U: UKCTOCS nested case-control study, A: The Avon Longitudinal Study of Children and Parents – mothers’ cohort, M: Metabolomics consortium, SD units: standard deviation units, CI: confidence interval.

**Supplementary table 1** – Estimated power in Mendelian randomization analysis

Exposure	Outcome	Sample size <sup>1</sup>	Type-I error rate	Effect estimate <sup>2</sup>	Instrument strength ( $R^2$ ) <sup>3</sup>	Power <sup>4</sup>
Adiponectin	Metabolic measure	23,884	0.05	0.05	0.04	34%
Adiponectin	Metabolic measure	23,884	0.05	0.10	0.04	87%
Adiponectin	Metabolic measure	23,884	0.05	0.20	0.04	88%

<sup>1</sup> Median sample size used for estimating SNP-outcome association

<sup>2</sup> Considering the true underlying causal association is unknown, a range of values was used.

<sup>3</sup> Instrument strength relates to the proportion of variance in the exposure explained by the instrument ( $R^2$ ). This was calculated by the sum of  $R^2$  from each 4 SNPs in the instrument. The formula used to estimate  $R^2$  for each SNP is detailed in Supplementary methods.

<sup>4</sup> We have estimated power for our Mendelian randomization analyses using the online calculator tool (<http://cnsgenomics.com/shiny/mRnd/>) and assuming a range of effect sizes for the potential underlying causal association between exposure and outcome.

**Supplementary table 2 – Heterogeneity estimates ( $I^2$ ) for meta-analysis of study-specific multivariate (MV) and Mendelian randomization (MR) estimates**

Metabolite	Overall		Females		Males		European studies only		Low risk individuals only	
	MV	MR	MV	MR	MV	MR	MV	MR	MV	MR
XXL-VLDL	85	0	82	7	NA	0	85	0	78	34
XL-VLDL	88	0	91	0	NA	0	88	0	79	0
L-VLDL	86	0	84	0	NA	11	86	0	82	0
M-VLDL	88	0	68	35	NA	52	88	0	83	0
S-VLDL	81	0	0	52	NA	63	81	0	79	19
XS-VLDL	0	23	0	73	NA	49	0	23	23	48
IDL	0	36	0	78	NA	44	0	36	23	57
L-LDL	0	43	0	78	NA	64	0	43	20	65
M-LDL	0	51	0	79	NA	63	0	51	32	71
S-LDL	0	53	0	78	NA	71	0	53	40	73
XL-HDL	66	26	88	0	NA	84	66	26	19	0
L-HDL	72	22	83	0	NA	81	72	22	59	28
M-HDL	0	57	0	0	NA	86	0	57	62	58
S-HDL	0	36	0	8	NA	63	0	36	4	50
VLDLd	83	0	26	0	NA	0	83	0	73	0
LDLd	0	25	0	0	NA	50	0	25	0	66
HDLd	75	18	64	32	NA	80	75	18	0	0
Total cholesterol	94	44	71	73	96	73	0	49	92	63
Remnant-c	87	23	0	73	NA	36	87	23	76	52
VLDL-c	89	4	0	61	NA	26	89	4	81	38
LDL-c	91	45	61	71	94	73	0	47	89	66
HDL-c	75	33	90	0	0	68	76	42	73	16
HDL2-c	74	57	70	0	NA	83	74	57	72	34
HDL3-c	0	45	93	43	NA	60	0	45	0	52
Esterified cholesterol	0	56	0	80	NA	77	0	56	0	68
Free cholesterol	43	46	31	77	NA	85	43	46	27	73
ApoA-I	0	55	79	0	NA	88	0	55	0	46
ApoB	86	20	0	73	NA	56	86	20	81	56
Triglycerides	73	0	91	39	48	50	80	0	70	0
Phosphoglycerides	0	48	58	48	NA	91	0	48	0	62
Phosphatidylcholines	0	58	41	62	NA	92	0	58	0	75
Total cholines	0	67	29	73	NA	90	0	67	0	74
Sphingomyelins	30	43	0	85	NA	0	30	43	0	45
DAG	NA	70	NA	0	NA	86	NA	70	NA	84
HDL-TG	0	47	75	44	NA	76	82	0	79	0
VLDL-TG	84	49	60	22	NA	36	84	49	68	40
LDL-TG	82	0	19	68	NA	78	0	47	42	56
TotFA	0	48	5	72	NA	86	0	48	42	68
SFA	0	45	0	65	NA	76	0	45	0	61
MUFA	91	43	91	61	NA	77	91	43	77	55
PUFA	32	64	45	79	NA	89	32	64	0	66
FAw6	10	42	35	77	NA	86	10	42	22	54
LA	0	0	16	70	NA	83	0	0	53	3
FAw3	47	79	13	82	NA	84	47	79	60	80
DHA	0	76	0	64	NA	89	0	76	16	80
CLA	NA	92	NA	0	NA	46	NA	92	NA	93
Glucose	0	54	46	0	57	49	21	51	34	39
HbA1c	62	0	11	0	NA	NA	NA	NA	21	38
Insulin	0	38	0	0	NA	53	0	38	37	0
Lactate	0	0	0	0	NA	0	0	0	0	0
Pyruvate	0	0	0	24	NA	NA	0	0	0	27
Citrate	69	0	0	29	NA	0	69	0	0	0
Glycerol	74	0	83	0	NA	NA	74	0	0	0
Alanine	65	0	62	0	NA	0	65	0	58	0
Glutamine	0	43	0	0	NA	75	0	43	0	58
Glycine	87	0	71	0	NA	NA	87	0	69	55
Isoleucine	0	0	0	0	NA	0	0	0	0	9
Leucine	0	34	0	0	NA	0	0	34	0	52
Valine	45	0	87	0	NA	0	45	0	0	0
Phenylalanine	32	72	30	0	NA	91	32	72	0	79
Tyrosine	0	48	40	0	NA	54	0	48	0	42
Histidine	0	10	0	0	NA	84	0	10	0	0
Acetoacetate	0	0	0	14	NA	NA	0	0	0	0
Acetate	0	31	0	35	NA	64	0	31	74	0
3-OH-butyrate	54	0	0	0	NA	0	54	0	0	0
Albumin	0	11	0	0	NA	0	0	11	0	26
Creatinine	0	61	0	49	NA	85	0	61	22	71
CRP	0	28	81	13	41	0	0	45	0	0
Fibrinogen	16	0	87	0	NA	60	16	0	0	17
IL-6	0	3	69	0	NA	57	0	3	0	0
GlycA	0	43	0	0	NA	39	0	43	58	0
Viscosity	0	56	0	0	NA	NA	0	56	0	0
SBP	0	0	28	11	44	0	0	0	0	0
DBP	68	0	72	36	0	0	83	0	0	0

NA: not applicable (estimates from only one study available). XXL: extremely large, XL: very large, L: large, M: medium, S: small, XS: very small, VLDL: very low-density lipoprotein, LDL: low-density lipoprotein, IDL: intermediate-density lipoprotein, HDL: high-density lipoprotein, c: cholesterol, DAG: diglycerides, TG: triglycerides, TotFA: total fatty acids, SFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acids, FAW6: omega-6 fatty acid, LA: linoleic acid, FAW3: omega-3 fatty acid, DHA: docosaenoic acid, CLA: conjugated linoleic acids, HbA1c: glycated haemoglobin, CRP: c-reactive protein, IL-6: interleukin-6, GlycA: glycoprotein acetyls, SBP: systolic blood pressure, DBP: diastolic blood pressure.



**Supplementary table 3.** P-values for the association of demographic and lifestyle variables with SNPs selected for Mendelian randomization analysis for each participating study

	PEL82	BWHHS	WHII	CaPS	UKCTOCS case- control*	ALSPAC- M
	<i>P-value</i>					
<i>Sex (male vs female)</i>						
rs6810075	0.15	—	0.67	—	—	—
rs16861209	0.12	—	0.45	—	—	—
rs17366568	0.36	—	0.84	—	—	—
rs3774261	0.35	—	0.63	—	—	—
<i>Age (years)</i>						
rs6810075	0.78	0.75	0.28	0.36	0.59	0.001
rs16861209	0.56	0.58	0.27	0.01	0.57	0.83
rs17366568	0.22	0.83	0.47	0.56	0.58	0.93
rs3774261	0.68	0.03	0.96	0.87	0.43	0.15
<i>European (yes vs no)</i>						
rs6810075	0.06	0.50	0.48	—	0.70	0.41
rs16861209	0.75	—	0.12	—	0.16	0.35
rs17366568	0.45	0.61	—	—	0.62	0.19
rs3774261	0.44	0.95	—	—	0.85	—
<i>Smoking (yes vs no)</i>						
rs6810075	0.64	0.22	0.77	0.48	—	0.11
rs16861209	0.57	0.37	0.87	0.48	—	0.24
rs17366568	0.45	0.62	0.44	0.77	—	0.90
rs3774261	0.52	0.08	0.90	0.37	—	0.92
<i>Body mass index (kg/m<sup>2</sup>)</i>						
rs6810075	0.63	0.49	0.49	0.21	0.05	0.45
rs16861209	0.39	0.41	0.59	0.65	0.20	0.72
rs17366568	0.47	0.73	0.87	0.11	0.66	0.32
rs3774261	0.48	0.65	0.41	0.44	0.78	0.04

ALSPAC-M: The Avon Longitudinal Study of Children and Parents – mothers' cohort; BWHHS: British Women's Heart and Health Study; CaPS: The Caerphilly Prospective Study; PEL82: 1982 Pelotas Birth Cohort; UKCTOCS: case-control study nested in The United Kingdom Collaborative Trial of Ovarian Cancer Screening; WHII: Whitehall-II Study.