

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

N-(3-oxododecanoyl)-L-homoserine lactone interactions in the breast tumor microenvironment: implications for breast cancer viability and proliferation *in vitro*

Brittany N. Balhouse^{1,2}, Logan Patterson^{3,4}, Eva M. Schmelz⁵, Daniel J. Slade³, Scott S. Verbridge^{1,2,*}

¹ School of Biomedical Engineering and Sciences, Virginia Tech-Wake Forest University, Blacksburg, VA, United States of America

² Department of Biomedical Engineering and Mechanics, Virginia Tech, Blacksburg, VA, United States of America

³ Department of Biochemistry, Virginia Tech, Blacksburg, VA, United States of America

⁴ Department of Pathology, University of Virginia, Charlottesville, VA, United States of America

⁵ Department of Human Nutrition, Foods and Exercise, Virginia Tech, Blacksburg, VA, United States of America

* Corresponding author

Email: sverb@vt.edu

25 **Abstract** It is well documented that the tumor microenvironment profoundly impacts the
26 etiology and progression of breast cancer, yet the contribution of the resident microbiome within
27 breast tissue remains poorly understood. Tumor microenvironmental conditions, such as hypoxia
28 and dense tumor stroma, predispose progressive phenotypes and therapy resistance, however the
29 role of bacteria in this interplay remains uncharacterized. We hypothesized that the effect of
30 individual bacterial secreted molecules on breast cancer viability and proliferation would be
31 modulated by these tumor-relevant stressors differentially for cells at varying stages of
32 progression. To test this, we incubated human breast adenocarcinoma cells (MDA-MB-231,
33 MCF-DCIS.com) and non-malignant breast epithelial cells (MCF-10A) with N-(3-
34 oxododecanoyl)-L-homoserine lactone (OdDHL), a quorum-sensing molecule from
35 *Pseudomonas aeruginosa* that regulates bacterial stress responses. This molecule was selected
36 because *Pseudomonas* was recently characterized as a significant fraction of the breast tissue
37 microbiome and OdDHL is documented to impact mammalian cell viability. After OdDHL
38 treatment, we demonstrated the greatest decrease in viability with the more malignant MDA-
39 MB-231 cells and an intermediate MCF-DCIS.com (ductal carcinoma *in situ*) response. The
40 responses were also culture condition (i.e. microenvironment) dependent. These results contrast
41 the MCF-10A response, which demonstrated no change in viability in any culture condition. We
42 further determined that the observed trends in breast cancer viability were due to modulation of
43 proliferation for both cell types, as well as the induction of necrosis for MDA-MB-231 cells in
44 all conditions. Our results provide evidence that bacterial quorum-sensing molecules interact
45 with the host tissue environment to modulate breast cancer viability and proliferation, and that
46 the effect of OdDHL is dependent on both cell type as well as microenvironment. Understanding
47 the interactions between bacterial signaling molecules and the host tissue environment will allow

48 for future studies that determine the contribution of bacteria to the onset, progression, and
49 therapy response of breast cancer.

50

51 **Introduction** The tumor microenvironment is now a widely recognized and well-studied
52 contributor to cancer dynamics, particularly for breast cancer. While increased matrix density,
53 programming of cancer-associated stromal cells, evolving gradients of oxygen and nutrients, and
54 leaky vasculature have all been implicated as key players in breast cancer progression (1-4), the
55 impact of the recently identified breast tissue resident microbiotic niche has received little
56 attention. Beyond the effects of pathogenic or tumorigenic bacteria such as *Chlamydomphila*
57 *pneumonia*, *Salmonella typhi*, *Streptococcus gallolyticus* (5), *Helicobacter pylori* (6) and
58 *Fusobacterium nucleatum* (7), the majority of analyses of tumor-microbiome interactions have
59 centered on local cell-cell interactions within the gut microenvironment, or more systemic
60 immune effects influenced by gut microbiota (8). Only a handful of studies have been conducted
61 to investigate the influences of tissue-resident bacteria in other tumor sites, such as for breast
62 cancer (9-11). Even fewer have investigated how small molecules released from resident bacteria
63 may interact with cells in the presence of other critical microenvironmental factors, e.g. tumor
64 hypoxia, to regulate cancer progression. In an effort to address these questions, we investigated
65 interactions between the quorum-sensing molecule N-(3-oxododecanoyl)-L-homoserine lactone
66 (OdDHL) and the breast tumor relevant microenvironmental cues of a stiff collagen-derived
67 tissue mimic and hypoxia. This representative study will aid in our understanding of how the
68 understudied breast tissue microbiome may contribute to disease phenotypes, patient-to-patient
69 variability, and cancer progression.

70

71 OdDHL secreting *Pseudomonas* are Gram-negative Proteobacteria that were recently found to
72 make up a significant fraction of the microbiome within breast tissue (12), and on nipple skin
73 and aspirate from women with and without a history of breast cancer (13). In addition,
74 *Pseudomonas* has been shown to make up a significant portion of the bacteria found in human
75 breast milk (11, 14-16). *Pseudomonas* was among the top five most abundant genera for both
76 sample populations in the 2014 study by Urbaniak et al. (12) and in several studies of breast milk
77 microbiota (14). OdDHL is a quorum-sensing molecule associated with biofilm development and
78 environmental stress response in bacteria (17) that has been shown to promote apoptosis in a
79 variety of human cell lines (18-23). Especially interesting is its selective effect inhibiting
80 proliferation and inducing apoptosis in breast cancer cells, but not in non-malignant breast cells
81 (21). For these reasons, studying the effects of OdDHL is not only important from the basic
82 science perspective (e.g. regarding its role in the tumor microenvironment), but also from the
83 therapeutic perspective given its potential as an anti-cancer treatment (24, 25).

84
85 High tissue density is not only a risk factor for breast cancer development, but also a clinical
86 diagnostic tool (26). Increased stroma density (resulting from increased collagen deposition and
87 cross-linking in the extracellular matrix (ECM)) is also associated with phenotypic changes in
88 both cancer cells and normal stroma cells (27). In an *in vitro* setting, growing cells on or within a
89 collagen hydrogel, as opposed to on a polystyrene substrate in two-dimensions (2D), changes not
90 only the type of focal adhesions cells make with their environment (28), but also how cells
91 respond to stresses within that environment (e.g. chemotherapeutic agents) (27, 29, 30). Three-
92 dimensional (3D) cellular adhesions are implicated in reduced cellular response to
93 chemotherapeutics (a phenomenon called adhesion-mediated resistance) and a progressive

94 phenotype (27, 31, 32). Hypoxia is also a well-documented constituent of the solid tumor
95 microenvironment; angiogenesis cannot keep up with the rate of tumor growth and produces a
96 gradient of oxygen from the well-fed periphery to a necrotic core that is devoid of oxygen (33).
97 Similar to cancer cells in a 3D environment, a low oxygen environment is associated with a
98 progressive phenotype and a resistance to chemotherapy (33, 34). Although, adhesion- and
99 hypoxia-mediated resistance have been extensively demonstrated for chemotherapeutic
100 compounds, adhesion- and hypoxia-mediated regulation of cellular response to bacterial
101 signaling molecules has not yet been investigated. Despite the previous research into OddHL,
102 the role of OddHL in the context of breast tumor microenvironmental stressors was still
103 unknown. Because such stressors are associated with altered cell responses (e.g.
104 chemoresistance), it was our hypothesis that OddHL would differentially modulate breast cancer
105 cell viability and proliferation in hypoxia and in 3D culture (in which cells are seeded in the bulk
106 of a tissue mimic).

107
108 In an effort to make a preliminary characterization of the response for both metastatic and non-
109 metastatic subtypes of breast cancer, we utilized MDA-MB-231 and MCF-DCIS.com cells,
110 respectively. MCF-10A breast epithelial cells were used as a non-malignant control. Both MDA-
111 MB-231 and MCF-10A cells were chosen for this research because of their wide documentation
112 in the literature (35, 36). The MCF-DCIS.com cell line, which is derived from the MCF-10A line
113 (37, 38), offers an intermediary model cell between the MCF-10A and MDA-MB-231 lines; not
114 only does it represent a pre-invasive, non-metastatic form of breast cancer, but is Her2+ (37)
115 whereas the MDA-MB-231 line is triple negative (35). In addition, both necrotic cores and
116 desmoplasia have been observed in lesions from MCF-DCIS.com injections (38) as well as in

117 MDA-MB-231 xenografts (39, 40), motivating our studies in hypoxia- and adhesion-mediated
118 resistance.

119

120 **Methods**

121 *Cell culture.* We used MDA-MB-231 triple negative breast cancer cells (ATCC® HTB-26™),
122 MCF-DCIS.com ductal carcinoma *in situ* breast cancer cells (generously donated by Dr. Eva M.
123 Schmelz) (38), and MCF-10A breast epithelial cells (ATCC® CRL-10317™) for our malignant
124 versus non-malignant cell comparison. MDA-MB-231 cells were cultured in DMEM/F12 (1:1)
125 media supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin (PS). MCF-
126 10A and MCF-DCIS.com cells were cultured in DMEM/F12 (1:1) media supplemented with 5%
127 horse serum, 1% PS, 0.05% hydrocortisone, 0.1% human insulin, 0.02% epidermal growth
128 factor, and 0.01% cholera toxin. Hereafter, these supplemented DMEM/F12 (1:1) media will be
129 referred to as complete media. Cells were subcultured every 4-5 days and used until passage 50.
130 Stock cells were grown in T-75 flasks under standard culture conditions (37°C incubator with
131 5% CO₂, and ambient O₂ (referred to hereafter as normoxia).

132

133 For 2D experimental conditions, cells were seeded in 48-well polystyrene plates at a
134 concentration of 100,000 cells per milliliter (30,000 cells per well), with two plates seeded per
135 cell type per experiment. After seeding, all plates were incubated under standard culture
136 conditions in normoxia. After 24 hours, the media was changed in all wells and the plates that
137 were to undergo the hypoxia treatment were moved into an incubator with 1% oxygen and the
138 standard culture conditions of 37°C, 5% CO₂.

139

140 *3D culture.* 3D culture was defined as cells suspended throughout a tissue mimic, namely a rat
141 tail derived type I collagen hydrogel. The collagen stock was manufactured using a technique
142 described previously (41). Briefly, collagen was extracted from Sprague-Dawley rat tails
143 (Bioreclamation, Inc), sterilized in 70% ethanol, and dissolved in 0.1% glacial acetic acid for
144 three days. The dissolved collagen stock was then aliquoted, frozen, lyophilized, and stored at –
145 20°C. At least three days prior to use, the lyophilized collagen was reconstituted in 0.1% glacial
146 acetic acid at a concentration of 12 milligrams of collagen per milliliter of acetic acid solution,
147 and this dissolved stock solution was then used for up to four weeks post re-constitution.

148
149 The hydrogel was created by neutralizing the collagen stock with empirically derived proportions
150 of 10X DMEM, 1X DMEM/F12 (1:1) and 1M sodium hydroxide (NaOH) (revised from (41)) to
151 create a hydrogel with pH 7.4 and 6 mg/mL of collagen in 0.1% acetic acid solution. During the
152 neutralization process, all solutions were kept on ice to prevent collagen polymerization. Cells
153 were suspended throughout the hydrogel by resuspension of a cell pellet into the required 1X
154 DMEM/F12 (1:1) prior to its addition into the unpolymerized (but neutralized) hydrogel
155 solution. Cells were seeded in the hydrogel solution at a concentration of three million cells per
156 milliliter of solution (~300,000 cells per well). The hydrogel solution was pipetted into
157 pretreated SYLGARD® 184 silicone elastomer molds topped with untreated coverslips to create
158 approximately 1 mm thick discs comprised of ~0.1 mL hydrogel solution. The hydrogels were
159 incubated for 30 minutes at 37°C, then transferred into 24-well plates and covered with 0.6 mL
160 complete media standard of the included cell type. There was one plate of 3D samples per cell
161 type. All 3D samples were maintained in an incubator under standard culture conditions (37°C,

162 5% CO₂, normoxia). Hydrogels with no cells were also created as a control for the alamarBlue™
163 cell viability assay described below.

164
165 *OddHL treatment.* Two days after seeding, the appropriate complete media containing varying
166 concentrations of OddHL were added to the samples. The concentrations of OddHL reflect
167 levels documented in the literature as physiological values in a niche in close proximity to the
168 bacteria producing the quorum sensing molecule (42). For the OddHL stock solution,
169 lyophilized OddHL (Sigma-Aldrich) was reconstituted in dimethyl sulfoxide (DMSO) at a
170 concentration of 67.25 mM or 20 mg/mL (the maximum solubility of OddHL in DMSO). For
171 the experimental solutions, the OddHL stock solution was added to the complete medium
172 appropriate for the cell line (as described above) to create the highest concentration (400 μM)
173 OddHL solution. Serial 1:2 dilutions of the stock solution were performed in DMSO to create
174 200 μM, 100 μM, 50 μM, and 25 μM solutions in complete medium. A control of 0.6% DMSO
175 (the volume per volume percentage of DMSO in all OddHL treatments) was used. Cells were
176 treated with the OddHL or control solutions for 24 hours before proliferation (by EdU assay)
177 and apoptosis/necrosis (by annexin V and propidium iodide assay) was analyzed, and 48 hours
178 before viability (by alamarBlue™ assay) was analyzed. The difference in incubation period was
179 determined by the nature of the assay to be performed.

180
181 *alamarBlue™ cell health assay.* After the OddHL or control treatment, the treatment solutions
182 were removed and fresh complete medium supplemented with 10% alamarBlue™ solution
183 (Thermo Fisher Scientific) was added to all wells, including 2D and 3D negative control wells
184 where there were no cells. These were incubated in normoxic or hypoxic incubators until

185 positive control wells (0.6% DMSO control) were magenta in color; reduction rate was
186 dependent on cell type and culture condition. When controls wells showed the color change,
187 three subsamples were taken from the solution of each well and added to a 96-well plate. As per
188 the manufacturer's instructions, the 96-well plates were read on a spectrophotometer, which
189 analyzed the absorbance at 570 and 600 nm wavelengths. The percent reduction of alamarBlue™
190 was calculated with the equation recommended by the manufacturer, namely:

$$\text{Percent Reduction of alamarBlue}^{\text{TM}} = \left(\frac{(O_2 \times A_1) - (O_1 \times A_2)}{(R_1 \times N_2) - (R_2 \times N_1)} \right) \times 100$$

191 where O_1 is the molar extinction coefficient (E) of oxidized alamarBlue™ at 570 nm, O_2 is the E
192 of oxidized alamarBlue™ at 600 nm, R_1 is the E of reduced alamarBlue™ at 570 nm, R_2 is the E
193 of reduced alamarBlue™ at 600 nm, A_1 is the absorbance of the test well at 570 nm, A_2 is the
194 absorbance of the test well at 600 nm, N_1 is the absorbance of the negative control at 570 nm and
195 N_2 is the absorbance of the negative control at 600 nm. The average percent reduction of
196 alamarBlue™ for each experimental group was divided by the average percent reduction of the 0
197 μM OdDHL (0.6% DMSO) control to calculate a relative viability.

198

199 *EdU assay.* An EdU assay was performed to study the effect of OdDHL on proliferation of cells
200 in normoxia (2D), hypoxia (2D), and 3D (normoxia). The EdU assay was conducted using the
201 Click-iT® EdU Alexa Fluor 488® Imaging Kit (Thermo Fisher Scientific) as per the
202 manufacturer's instructions, with the exception of the fixing where 10% formalin was substituted
203 for 3.7% formaldehyde. Briefly, after incubation with OdDHL in the various experimental
204 conditions for 24 hours, half of the experimental solution was removed and fresh complete
205 medium supplemented with the EdU molecule was added to each well (including controls). After
206 a two hour incubation, the samples were fixed with 10% formalin (15 minutes for 2D, 45

207 minutes for 3D), washed and incubated with the Click-iT® reaction cocktail to fluorescently
208 label cells that entered the S-phase of the cell cycle during the two hour incubation period. Cells
209 were then counterstained with DAPI. Wells were imaged under 100X magnification with 1
210 subsample per well for 2D conditions and, using a confocal microscope, 5 subsamples per well
211 for 3D conditions vertically through the hydrogel volume (30 µm spacing). The images were
212 post-processed in ImageJ; see Supporting Information for more information on post-processing
213 (Fig S1). The area of cell nuclei stained with Alexa Fluor 488® was divided by the area of cell
214 nuclei stained with DAPI to find the fraction of cells entering S-phase during the 2 hour
215 incubation period.

216

217 *Apoptosis/necrosis assay.* The apoptosis and necrosis response was determined by incubation
218 with annexin V-FITC (AV-FITC) and propidium iodide (PI) using a commercial assay (Annexin
219 V-FITC Apoptosis Kit, BioVision Inc.). The assay was completed according to manufacturer
220 instructions for all 2D samples and with three times the recommended incubation time for 3D
221 samples. After incubation, cells were immediately imaged using 100X magnification with 3
222 subsamples per well for 2D conditions and, using a confocal microscope, 5 subsamples per well
223 for 3D conditions vertically through the hydrogel volume (30 µm spacing). The images were
224 post-processed in ImageJ; see Supporting Information for more information on post-processing
225 (Fig S2). All values are relative to the appropriate 0.6% DMSO control for each experimental
226 condition.

227

228 *Statistical analysis.* For each experiment, there were two wells per experimental condition and
229 control and each experiment was conducted three times ($N = 3$), unless otherwise noted. The
230 statistics are based on the average values for each condition in each experiment.

231
232 The statistical analyses consisted of a full factorial two-way analysis of variance (ANOVA) in
233 which the differences in viability (relative percent reduction of alamarBlue™), proliferation rate
234 (percent of cells entering S-phase), and apoptosis/necrosis (AV-FITC/PI stained cells) for
235 varying culture conditions and OddHL concentrations were analyzed for each cell line tested.
236 Where a p-value < 0.05 was found for the ANOVA, post-hoc Tukey tests and least means
237 contrasts were performed to analyze the degree of significance between experimental conditions.
238 Error bars represent standard error of the mean.

239
240 **Results** We first compared highly malignant MDA-MB-231 cells with non-malignant MCF-10A
241 cells. We found that the response to OddHL was dependent not only on cell-type, but also the
242 culture condition. As compared to the control, the malignant MDA-MB-231 cells showed
243 significantly different responses to 400 μM of OddHL in all culture conditions (Fig 1A-1E). In
244 the 2D/normoxia condition, the 400 μM OddHL treatment corresponded to approximately
245 52.2% ($\pm 2.2\%$) viability, relative to the control. In the 2D/hypoxia condition, that relative
246 viability was increased to 60.6% ($\pm 2.2\%$) and in 3D/normoxia condition, that viability was
247 increased to 81.9% ($\pm 2.2\%$) (Fig 1C and Fig 1D, respectively). There were significant decreases
248 in the MDA-MB-231 viability at the 100 and 200 μM levels of OddHL, as compared to the
249 control, for the 2D conditions as well (Fig 1A and Fig 1B). In contrast, the non-malignant MCF-

250 MCF-10A cells exposed to OddHL showed no significant change in viability relative to the control
251 across all concentrations in all culture conditions (Fig 1F-1H).

252

253 **Figure 1. OddHL has a cell-specific and significant impact on MDA-MB-231 viability in all**
254 **culture conditions.**

255 Significant differences in mean MDA-MB-231 viability at 100, 200 and 400 μ M concentrations
256 relative to the control in hypoxia and normoxia in 2D (A) and 3D and 2D in normoxia (B); *** =
257 p-value < 0.0001, ** = p-value < 0.01, * = p-value < 0.05 based on Tukey post-hoc differences.

258 Significant differences in MDA-MB-231 viability relative to the control between culture
259 conditions with 400 μ M treatment in hypoxia and normoxia in 2D (C) and 3D and 2D in
260 normoxia (D); *** = p-value < 0.0001, ** = p-value < 0.01 based on least mean contrasts. (E)

261 Photomicrographs of MDA-MB-231 cells cultured in normoxia (2D), hypoxia (2D), and 3D
262 (normoxia) with fluorescence microscopy (with DAPI staining; scale bar = 100 μ m) at 100X

263 magnification after 24 hours of OddHL treatment. (F) Photomicrographs of MCF-10A cells
264 cultured in normoxia (2D), hypoxia (2D), and 3D (normoxia) with fluorescence microscopy

265 (with DAPI staining; scale bar = 100 μ m) at 100X magnification after 24 hours of OddHL
266 treatment. Differences in MCF-10A viability relative to the control at 100, 200 and 400 μ M

267 concentrations in hypoxia and normoxia in 2D (G) and 3D and 2D in normoxia (H). Error bars
268 represent standard error of the mean.

269

270 We next sought to quantify the viability response of the MCF-DCIS.com cells, which represent
271 an intermediate phenotype between the non-malignant and metastatic extremes. The viability
272 assay of the MCF-DCIS.com cells showed the cells responded in a similar fashion to MDA-MB-

273 231 cells in both the 2D/normoxia and 2D/hypoxia conditions, with significant dose-dependent
274 decreases in viability with increasing OdDHL concentrations (Fig 2A). However, unlike the
275 MDA-MB-231 cells, the MCF-DCIS.com cells did not have a significant decrease in viability in
276 the 3D/normoxia condition at any treatment level of OdDHL (Fig 2B); this is an interesting
277 observation to which we will return shortly. Also, unique to the MCF-DCIS.com cells is a
278 significantly lower viability at the 400 μ M level for MCF-DCIS.com cells cultured in
279 2D/hypoxia than those cultured in 2D/normoxia (Fig 2C). However, for MCF-DCIS.com cells
280 grown in 3D versus 2D in normoxia, there was significantly increased MCF-DCIS.com viability
281 with the 400 μ M OdDHL treatment (Fig 2D). An analysis of the cell specific viability response
282 to 400 μ M OdDHL showed that MCF-DCIS.com cells have significantly higher viability than
283 MDA-MB-231 cells, but significantly lower viability than MCF-10A cells when cultured under
284 the 2D/normoxia condition (Fig 2E). With the same treatment, MCF-DCIS.com cells showed
285 significantly lower viability than MCF-10A cells when cultured under the 2D/hypoxia (a
286 response that is statistically not different from that of MDA-MB-231 cells under the same
287 condition) (Fig 2E). When cells were cultured under the 3D/normoxia condition, there was no
288 difference in MCF-DCIS.com viability as compared to MCF-10A cell viability with the 400 μ M
289 OdDHL treatment; both were significantly higher than MDA-MB-231 cell viability under the
290 same condition (Fig 2F).

291
292 **Figure 2. OdDHL has a cell-specific and differential impact on MCF-DCIS.com cells**
293 **dependent on culture condition.**

294 Significant differences in mean MCF-DCIS.com viability relative to the control at 100, 200 and
295 400 μ M OdDHL concentrations in hypoxia and normoxia in 2D (A) and 3D and 2D in normoxia

296 (B); *** = p-value < 0.0001, ** = p-value < 0.01, * = p-value < 0.05 based on Tukey post-hoc
297 differences. Significant differences in MCF-DCIS.com viability relative to the control between
298 culture conditions with 400 μ M OddHL treatment in hypoxia and normoxia in 2D (C) and 3D
299 and 2D in normoxia (D); *** = p-value < 0.0001 based on least mean contrasts. Significant
300 differences in mean MCF-10A, MCF-DCIS.com, MDA-MB-231 viabilities relative to the
301 control at 400 μ M OddHL concentrations in hypoxia and normoxia in 2D (E) and 3D and 2D in
302 normoxia (F); *** = p-value < 0.0001 based on Tukey post-hoc differences. Error bars represent
303 standard error of the mean.

304
305 To provide further clarity regarding the variations in viability that we observed in different
306 culture conditions for the malignant cells (MDA-MB-231 and MCF-DCIS.com), we next
307 quantified proliferation by measuring S-phase entry. The results of the proliferation assay
308 paralleled the results of the viability assay for both the MDA-MB-231 and MCF-DCIS.com
309 cells. The percentage of cells entering the S-phase of the cell cycle during the two-hour
310 incubation period with EdU was significantly lower for 400 μ M OddHL-treated MDA-MB-231
311 cells than for control (0.6% DMSO) cells across all culture conditions (Fig 3A, Fig 3B). The
312 percentage of EdU-tagged cells after the two hour incubation period was also lower for 400 μ M
313 OddHL-treated versus the control cells across all culture conditions for the MCF-DCIS.com
314 cells (Fig 3C, Fig 3D).

315
316 **Figure 3. 400 μ M OddHL treatment significantly impacts the proliferation rate of MDA-**
317 **MB-231 and MCF-DCIS.com cells in all conditions.**

318 Significant differences in the mean percent of MDA-MB-231 cells entering S-phase of
319 proliferation in the 0.6% DMSO and 400 μ M OdDHL treatment groups in hypoxia and normoxia
320 in 2D for N = 4 (A) and 3D and 2D in normoxia for N = 3 (B); *** = p-value < 0.0001 based on
321 least mean contrasts. Significant differences in the mean percent of MCF-DCIS.com cells
322 entering S-phase of proliferation in the 0.6% DMSO and 400 μ M OdDHL treatment groups in
323 hypoxia and normoxia in 2D for N = 3 (C) and 3D and 2D in normoxia for N = 3 (D); ** = p-
324 value < 0.001, * = p-value < 0.05 based on least mean contrasts. Error bars represent standard
325 error of the mean.

326
327 To provide further insights into the viability data, especially in any case where the viability and
328 proliferation trends were not consistent (i.e. the 3D/normoxia condition for MCF-DCIS.com
329 cells), we next quantified apoptosis and necrosis for all cell types. Not surprisingly, the
330 apoptosis/necrosis assay revealed that OdDHL generally induces cell death exclusively for
331 malignant cells. Analysis of the mean gray value (MGV) for the AV-FITC staining showed that
332 apoptosis of MDA-MB-231 cells with 400 μ M OdDHL relative to the 0.6% DMSO control in all
333 culture conditions was not significantly increased (Fig 4A). Analysis of the MGV for the PI
334 staining showed significantly increased necrosis of MDA-MB-231 with 400 μ M OdDHL
335 treatment, relative to the 0.6% DMSO control, in all culture conditions (Fig 4B). Analysis of
336 MGV for the AV-FITC and PI staining of MCF-DCIS.com cells showed significantly increased
337 apoptosis and necrosis, respectively, only for cells in 2D/normoxia with the 400 μ M OdDHL
338 treatment (Fig 4C and Fig 4D). The PI staining of the MCF-DCIS.com cells in the 3D/normoxia
339 showed that these cells had decreased necrosis with 400 μ M OdDHL treatment relative to the
340 control, a point to which we will return in the discussion (Fig 4D). The 400 μ M OdDHL did not

341 significantly increase apoptosis or necrosis of MCF-10A cells compared to the 0.6% DMSO
342 control (Fig 4E and Fig 4F). Rather, 400 μ M OddHL significantly decreased both MCF-10A
343 apoptosis and necrosis under the 3D/normoxia condition compared to the 0.6% DMSO control
344 (Fig 4E and Fig 4F).

345

346 **Figure 4. 400 μ M OddHL treatment significantly increases necrosis of MDA-MB-231 cells**
347 **in all conditions.**

348 (A) Differences in the annexin V-FITC staining mean gray value (MGV) for MDA-MB-231
349 cells treated with 400 μ M OddHL, normalized to 0.6% DMSO control for 3D/normoxia,
350 2D/hypoxia, and 2D/normoxia culture conditions (N = 3). (B) Differences in the propidium
351 iodide staining MGV for MDA-MB-231 cells treated with 400 μ M OddHL, normalized to 0.6%
352 DMSO control for 3D/normoxia, 2D/hypoxia, and 2D/normoxia culture conditions (N = 3); ** =
353 p-value < 0.01, * = p-value < 0.05 based on least mean contrasts. (C) Differences in the annexin
354 V-FITC staining MGV for MCF-DCIS.com cells treated with 400 μ M OddHL, normalized to
355 0.6% DMSO control for 3D/normoxia (N = 3), 2D/hypoxia (N = 4), and 2D/normoxia culture
356 conditions (N = 4); *** = p-value < 0.001 based on least mean contrasts. (D) Differences in the
357 propidium iodide staining MGV for MCF-DCIS.com cells treated with 400 μ M OddHL,
358 normalized to 0.6% DMSO control for 3D/normoxia, 2D/hypoxia, and 2D/normoxia culture
359 conditions (N = 3); ** = p-value < 0.01, * = p-value < 0.05 based on least mean contrasts. (E)
360 Differences in the annexin V-FITC staining MGV for MCF-10A cells treated with 400 μ M
361 OddHL, normalized to 0.6% DMSO control for 3D/normoxia, 2D/hypoxia, and 2D/normoxia
362 culture conditions (N = 3); ** = p-value < 0.01 based on least mean contrasts. (F) Differences in
363 the propidium iodide staining MGV for MCF-10A cells treated with 400 μ M OddHL,

364 normalized to 0.6% DMSO control for 3D/normoxia, 2D/hypoxia, and 2D/normoxia culture
365 conditions (N = 3); ** = p-value < 0.01 based on least mean contrasts. Error bars represent
366 standard error of the mean.

367
368 **Discussion** The analysis of the viability of MDA-MB-231 and MCF-10A cells in response to
369 OdDHL confirmed our hypotheses that there would be reduced MDA-MB-231 response to
370 OdDHL treatment in both hypoxia and in 3D while MCF-10A cells would show no such
371 changes. Our 2D/normoxia results are consistent with published literature (21), while our other
372 conditions provide further insight into the interaction of OdDHL with other tumor
373 microenvironment factors. As we anticipated, malignant cells cultured in a 3D environment had
374 the highest relative viability (Fig 1) with OdDHL treatment as compared to cells in 2D. Cells in
375 hypoxia also had increased relative viability with OdDHL treatment as compared to those in
376 normoxia (Fig 1). Thus, we found that OdDHL preferentially affects MDA-MB-231 cells as
377 compared to MCF-10A cells and this effect is blunted by both hypoxia and 3D culture. This
378 result parallels the common finding that tumor cells are more resistant to chemotherapies when
379 studied in 3D and/or hypoxic conditions (27, 31-34), and suggests that these conditions also
380 represent a more physiologically relevant context for the testing of the impact of microbial
381 factors in the tumor microenvironment.

382
383 It is interesting to note that, while MDA-MB-231 cells had a decreased viability response when
384 cultured under hypoxia and 3D conditions, the intermediate malignancy MCF-DCIS.com cells
385 did not respond in the same way (Fig 2). MCF-DCIS.com viability was in-between that of the
386 MCF-10A and MDA-MB-231 cells in 2D/normoxia, statistically similar to the MDA-MB-231

387 response in 2D/hypoxia, and statistically similar to the MCF-10A response in 3D/normoxia (Fig
388 2). This novel evidence suggests that bacterial factors may affect mammalian cells differently at
389 each stage of cancer progression. It is possible that the response to OddHL is linked to
390 epithelial-to-mesenchymal transition as it has previously been shown that OddHL has an effect
391 on cytoskeletal proteins in multiple malignant cell lines (24, 43). OddHL may be differentially
392 affecting the MDA-MB-231 cells in comparison to the MCF-DCIS.com cells due to its
393 mesenchymal and highly metastatic characteristics; however, much more work needs to be done
394 for confirmation of this hypothesis.

395
396 In order to better understand the changes in viability for the breast cancer cells as measured by
397 the alamarBlue™ assay, we next decided to examine the effect of OddHL on both cell
398 proliferation and apoptosis/necrosis. For MDA-MB-231 cells, it was revealed that OddHL has
399 two mechanisms to decrease the cell number. We found that OddHL significantly decreases
400 proliferation of MDA-MB-231 cells in all culture conditions (Fig 3). We then hypothesized that
401 decrease in proliferation alone could not account for the decrease in viability we observed. A
402 second mechanism of action was found in the apoptosis/necrosis assay. We found that necrosis
403 of MDA-MB-231 cells was significantly increased with the 400 μ M OddHL treatment as
404 compared to the 0.6% DMSO control (Fig 4). These two factors worked in concert resulting in
405 the significant viability decrease observed for MDA-MB-231 cells in all conditions.

406
407 These effects were not seen in MCF-10A cells and, in fact, both apoptosis and necrosis were
408 reduced in the 3D/normoxia condition while remaining unchanged in all other conditions (Fig 4).
409 The fact that OddHL triggers necrosis as opposed to apoptosis in malignant cells is not

410 surprising, as bacteria are known to induce necrosis (44). However, the fact that OddHL may
411 reduce apoptosis and necrosis for non-malignant cells under a stressful microenvironment (i.e. a
412 stiff collagen matrix) is intriguing. More research is needed to understand the selectivity of
413 necrosis induction and, especially, reduction by OddHL.

414
415 Likewise, we found that the primary mechanism responsible for the decrease in MCF-DCIS.com
416 cell number seen with OddHL treatment was decreased proliferation (Fig 3). While we did not
417 observe decreased viability for MCF-DCIS.com cells in the 3D/normoxia condition, decreased
418 proliferation was observed in all three culture conditions. The apoptosis/necrosis assay shed
419 some light on this seemingly contradictory result. It was found that, as with the MCF-10A cells,
420 there was decreased necrosis for MCF-DCIS.com cells in the 3D/normoxia condition (Fig 4).
421 From these data, we propose that the opposing effects of the decreased proliferation and the
422 decreased necrosis caused by the 400 μ M OddHL treatment countered each other, leading to the
423 invariable response in viability observed. The MCF-DCIS.com cells exhibited increased
424 apoptosis and necrosis in the 2D/normoxia condition, which would have worked in concert with
425 decreased proliferation in this condition leading to the reduced viability observed (Fig 4).
426 Interestingly, we note that the largest reduction in viability for MCF-DCIS.com cells treated with
427 OddHL was in the 2D/hypoxia condition, although this condition did not result in any
428 significant increases in apoptosis or necrosis. This is different than the observation for the highly
429 malignant MDA-MB-231 cells, whose viability decreased the most under 2D/normoxia. While
430 this may relate to the fact that hypoxia is linked to and predisposes a highly malignant phenotype
431 (45, 46), future mechanistic work will be needed to further clarify these observations. Taken
432 together, these data are indicative that there may be different mechanisms of OddHL action on

433 cancer viability and these are cell type and microenvironmental condition dependent. While
434 more work with cells representative of varying stages of progression needs to be completed,
435 these preliminary data suggest that OddHL may be more effective at suppressing the growth of
436 more malignant or advanced breast cancer.

437
438 Future studies investigating the possible role of tumor microenvironmental microbiota should
439 establish if, as is suggested here, OddHL suppresses highly malignant phenotypes. In addition,
440 further experiments should extend the study of non-malignant and pre-malignant cell types to
441 establish if, as seems to be the case for MCF-10A cells, OddHL increases the survival of these
442 cells in the presence of tumor microenvironmental stressors. Finally, although we show that
443 OddHL decreases breast viability through inhibition of proliferation as well as cell type and
444 culture condition dependent induction of apoptosis and/or necrosis, future work should establish
445 the cell signaling pathways associated with the differential responses to OddHL. Li et al. linked
446 the selective action of OddHL on malignant cell lines to STAT3 activity (21). However, further
447 studies into the mechanism of action of OddHL with our experimental conditions are required to
448 see if this explanation also extends to our data.

449
450 The breast cancer response to OddHL described above demonstrates that there is significant
451 interaction between bacterial factors and chemical or physical stresses (e.g. hypoxia and stiff
452 ECM) in the tumor microenvironment. Interestingly, in the case of MDA-MB-231 and MCF-
453 DCIS.com cells, we demonstrated that malignant cell hypoxia- and adhesion-mediated resistance
454 are not purely associated with chemotherapeutic compounds, but with native microbiome
455 signaling molecules as well. However, it is also notable that, although the MDA-MB-231 cells

456 demonstrated a blunted response to OddHL in hypoxia and in 3D, there was still a significant
457 response. Thus, future studies into both microbiome-tumor interactions and the therapeutic
458 potential of OddHL could provide valuable information.

459
460 Many open questions remain regarding the physiological or pathological role of the breast
461 resident microbiome. Prior research has established an association between the composition of
462 the breast microbiota and cancer, demonstrating the ability of the abundant bacteria in breast
463 cancer patients to induce double-stranded DNA breaks and/or cell proliferation (9). However, it
464 is not yet known whether these microbiome compositions are a cause or consequence of the
465 malignancies (9). Our data provide an important contribution to this field, demonstrating that
466 even for one quorum-sensing molecule there is a complex breast cell response that is dependent
467 on both cell type as well as microenvironmental context. Many more studies of breast relevant
468 bacteria secreted molecules are needed to establish the impact of the soluble factor crosstalk
469 between the microbiome and breast cancer during the entire course of progression as well as
470 during therapy response. With our data, we have established the importance of performing these
471 investigations in the context of physiologically relevant *in vitro* models. Thus, such
472 investigations would benefit from the incorporation of microenvironmental stressors. More
473 broadly, co-culture experiments investigating soluble factor interactions between breast tissue
474 relevant bacteria and cancer cells in the context of tumor microenvironmental stressors could
475 provide valuable insights into carcinogenesis as well as patient-to-patient variability in disease
476 course and therapy response.

477

478 **Conclusions** We showed that the selective effect of OddHL on the viability of breast cancer
479 cells is significantly mitigated by two key hallmarks of the tumor microenvironment: hypoxia
480 and stiff ECM. It is important to note that, although there was a blunted response to OddHL
481 treatment for MDA-MB-231 cells in hypoxia and in 3D, there was still a significant decrease in
482 proliferation and viability, and a significant increase in necrosis with treatment. The intermediary
483 response of MCF-DCIS.com may demonstrate a differential role of microenvironmental bacterial
484 factors over the course of breast cancer progression. Thus, further exploration of microbiome-
485 tumor interactions and OddHL as a potential therapy is merited.

486

487 **Acknowledgements** We would like to thank Sara Peterson for her assistance in experimental
488 preparation. We would also like to thank Dr. Akanksha Kanitkar for her thoughtful discussions,
489 and Dr. Ann Stevens for useful input on the role of OddHL in bacterial stress response.

490

491 Works Cited

492

493 1. Shieh AC. Biomechanical forces shape the tumor microenvironment. *Annals of*
494 *biomedical engineering*. 2011;39(5):1379-89.

495 2. Koumoutsakos P, Pivkin I, Milde F. *The Fluid Mechanics of Cancer and Its Therapy*.
496 *Annu Rev Fluid Mech*. 2013;45:325-55.

497 3. Jain RK. Normalizing tumor microenvironment to treat cancer: bench to bedside to
498 biomarkers. *Journal of clinical oncology : official journal of the American Society of Clinical*
499 *Oncology*. 2013;31(17):2205-18.

500 4. Quail DF, Joyce JA. Microenvironmental regulation of tumor progression and metastasis.
501 *Nature medicine*. 2013;19(11):1423-37.

502 5. Cummins J, Tangney M. Bacteria and tumours: causative agents or opportunistic
503 inhabitants? *Infectious agents and cancer*. 2013;8(1):1.

504 6. Polk DB, Peek RM. *Helicobacter pylori*: gastric cancer and beyond. *Nature Reviews*
505 *Cancer*. 2010;10(6):403-14.

506 7. Hieken TJ, Chen J, Hoskin TL, Walther-Antonio M, Johnson S, Ramaker S, et al. The
507 Microbiome of Aseptically Collected Human Breast Tissue in Benign and Malignant Disease.
508 *Scientific reports*. 2016;6:30751.

509 8. Garrett WS. Cancer and the microbiota. *Science*. 2015;348(6230):80-6.

510 9. Urbaniak C, Gloor GB, Brackstone M, Scott L, Tangney M, Reid G. The microbiota of
511 breast tissue and its association with breast cancer. *Applied and environmental microbiology*.
512 2016;82(16):5039-48.

- 513 10. Xuan C, Shamonki JM, Chung A, DiNome ML, Chung M, Sieling PA, et al. Microbial
514 dysbiosis is associated with human breast cancer. *PloS one*. 2014;9(1):e83744.
- 515 11. Reid G. Can breast microbiota provide protective effects against cancer? : *Future*
516 *Medicine*; 2016.
- 517 12. Urbaniak C, Cummins J, Brackstone M, Macklaim JM, Gloor GB, Baban CK, et al.
518 Microbiota of human breast tissue. *Applied and environmental microbiology*. 2014;80(10):3007-
519 14.
- 520 13. Chan AA, Bashir M, Rivas MN, Duvall K, Sieling PA, Pieber TR, et al. Characterization
521 of the microbiome of nipple aspirate fluid of breast cancer survivors. *Scientific reports*.
522 2016;6:28061.
- 523 14. McGuire MK, McGuire MA. Got bacteria? The astounding, yet not-so-surprising,
524 microbiome of human milk. *Current Opinion in Biotechnology*. 2017;44:63-8.
- 525 15. Gomez-Gallego C, Garcia-Mantrana I, Salminen S, Collado MC. The human milk
526 microbiome and factors influencing its composition and activity. *Seminars in Fetal and Neonatal*
527 *Medicine*. 2016;21(6):400-5.
- 528 16. Hoashi M, Meche L, Mahal LK, Bakacs E, Nardella D, Naftolin F, et al. Human Milk
529 Bacterial and Glycosylation Patterns Differ by Delivery Mode. *Reproductive Sciences*.
530 2015;23(7):902-7.
- 531 17. García-Contreras R, Nunez-Lopez L, Jasso-Chávez R, Kwan BW, Belmont JA, Rangel-
532 Vega A, et al. Quorum sensing enhancement of the stress response promotes resistance to
533 quorum quenching and prevents social cheating. *The ISME journal*. 2015;9(1):115-25.

- 534 18. Tao S, Luo Y, He B, Liu J, Qian X, Ni Y, et al. Paraoxonase 2 modulates a proapoptotic
535 function in LS174T cells in response to quorum sensing molecule N-(3-oxododecanoyl)-L-
536 homoserine lactone. *Scientific reports*. 2016;6.
- 537 19. Schwarzer C, Fu Z, Patanwala M, Hum L, Lopez-Guzman M, Illek B, et al. *Pseudomonas*
538 *aeruginosa* biofilm-associated homoserine lactone C12 rapidly activates apoptosis in airway
539 epithelia. *Cellular microbiology*. 2012;14(5):698-709.
- 540 20. Tateda K, Ishii Y, Horikawa M, Matsumoto T, Miyairi S, Pechere JC, et al. The
541 *Pseudomonas aeruginosa* Autoinducer N-3-Oxododecanoyl Homoserine Lactone Accelerates
542 Apoptosis in Macrophages and Neutrophils. *Infection and Immunity*. 2003;71(10):5785-93.
- 543 21. Li L, Hooi D, Chhabra SR, Pritchard D, Shaw PE. Bacterial N-acylhomoserine lactone-
544 induced apoptosis in breast carcinoma cells correlated with down-modulation of STAT3.
545 *Oncogene*. 2004;23(28):4894-902.
- 546 22. Shiner E, Terentyev D, Bryan A, Sennoune S, Martinez-Zaguilan R, Li G, et al.
547 *Pseudomonas aeruginosa* autoinducer modulates host cell responses through calcium signalling.
548 *Cellular microbiology*. 2006;8(10):1601-10.
- 549 23. Jacobi CA, Schiffner F, Henkel M, Waibel M, Stork B, Daubrawa M, et al. Effects of
550 bacterial N-acyl homoserine lactones on human Jurkat T lymphocytes-OddHL induces apoptosis
551 via the mitochondrial pathway. *International Journal of Medical Microbiology*. 2009;299(7):509-
552 19.
- 553 24. Kumar AS, Bryan JN, Kumar SR. Bacterial quorum sensing molecule N-3-oxo-
554 dodecanoyl-L-homoserine lactone causes direct cytotoxicity and reduced cell motility in human
555 pancreatic carcinoma cells. *PloS one*. 2014;9(9):e106480.

- 556 25. Zhao G, Neely AM, Schwarzer C, Lu H, Whitt AG, Stivers NS, et al. N-(3-oxo-acyl)
557 homoserine lactone inhibits tumor growth independent of Bcl-2 proteins. *Oncotarget*.
558 2016;7(5):5924-42.
- 559 26. Checka CM, Chun JE, Schnabel FR, Lee J, Toth H. The relationship of mammographic
560 density and age: implications for breast cancer screening. *American Journal of Roentgenology*.
561 2012;198(3):W292-W5.
- 562 27. Infanger DW, Lynch ME, Fischbach C. Engineered culture models for studies of tumor-
563 microenvironment interactions. *Annual review of biomedical engineering*. 2013;15:29-53.
- 564 28. Pelham RJ, Wang Y-l. Cell locomotion and focal adhesions are regulated by substrate
565 flexibility. *Proceedings of the National Academy of Sciences*. 1997;94(25):13661-5.
- 566 29. Edmondson R, Broglie JJ, Adcock AF, Yang L. Three-dimensional cell culture systems
567 and their applications in drug discovery and cell-based biosensors. *Assay and drug development*
568 *technologies*. 2014;12(4):207-18.
- 569 30. DelNero P, Lane M, Verbridge SS, Kwee B, Kermani P, Hempstead B, et al. 3D culture
570 broadly regulates tumor cell hypoxia response and angiogenesis via pro-inflammatory pathways.
571 *Biomaterials*. 2015;55:110-8.
- 572 31. Seguin L, Desgrosellier JS, Weis SM, Cheresch DA. Integrins and cancer: regulators of
573 cancer stemness, metastasis, and drug resistance. *Trends in cell biology*. 2015;25(4):234-40.
- 574 32. Eke I, Cordes N, editors. *Focal adhesion signaling and therapy resistance in cancer*.
575 *Seminars in cancer biology*; 2015: Elsevier.
- 576 33. Ward C, Langdon SP, Mullen P, Harris AL, Harrison DJ, Supuran CT, et al. New
577 strategies for targeting the hypoxic tumour microenvironment in breast cancer. *Cancer treatment*
578 *reviews*. 2013;39(2):171-9.

- 579 34. Rohwer N, Cramer T. Hypoxia-mediated drug resistance: novel insights on the functional
580 interaction of HIFs and cell death pathways. *Drug Resistance Updates*. 2011;14(3):191-201.
- 581 35. Chavez KJ, Garimella SV, Lipkowitz S. Triple Negative Breast Cancer Cell Lines: One
582 Tool in the Search for Better Treatment of Triple Negative Breast Cancer. *Breast disease*.
583 2010;32(1-2):35-48.
- 584 36. Cowell JK, LaDuca J, Rossi MR, Burkhardt T, Nowak NJ, Matsui S-i. Molecular
585 characterization of the t(3;9) associated with immortalization in the MCF10A cell line. *Cancer*
586 *Genetics and Cytogenetics*. 2005;163(1):23-9.
- 587 37. So JY, Lee HJ, Kramata P, Minden A, Suh N. Differential Expression of Key Signaling
588 Proteins in MCF10 Cell Lines, a Human Breast Cancer Progression Model. *Molecular and*
589 *cellular pharmacology*. 2012;4(1):31-40.
- 590 38. Miller FR, Santner SJ, Tait L, Dawson PJ. MCF10DCIS. com xenograft model of human
591 comedo ductal carcinoma in situ. *Journal of the National Cancer Institute*. 2000;92(14):1185a-6.
- 592 39. Yen T-H, Lee G-D, Chai J-W, Liao J-W, Lau J-Y, Hu L-C, et al. Characterization of a
593 murine xenograft model for contrast agent development in breast lesion malignancy assessment.
594 *Journal of Biomedical Science*. 2016;23(1):46.
- 595 40. Nelson MT, Short A, Cole SL, Gross AC, Winter J, Eubank TD, et al. Preferential,
596 enhanced breast cancer cell migration on biomimetic electrospun nanofiber ‘cell highways’.
597 *BMC cancer*. 2014;14(1):825.
- 598 41. Cross VL, Zheng Y, Choi NW, Verbridge SS, Sutermaster BA, Bonassar LJ, et al. Dense
599 type I collagen matrices that support cellular remodeling and microfabrication for studies of
600 tumor angiogenesis and vasculogenesis in vitro. *Biomaterials*. 2010;31(33):8596-607.

- 601 42. Lazenby JJ. The role of N-(3-oxododecanoyl)-L-homoserine lactone in the pathogenesis
602 of *Pseudomonas aeruginosa* [Ph.D.]: University of New South Wales; 2011.
- 603 43. Dolnick R, Wu Q, Angelino NJ, Stephanie LV, Chow K-C, Sufrin JR, et al. Enhancement
604 of 5-Fluorouracil Sensitivity by an rTS Signaling Mimic in H630 Colon Cancer Cells. *Cancer*
605 *Research*. 2005;65(13):5917-24.
- 606 44. Proskuryakov SYa, Konoplyannikov AG, Gabai VL. Necrosis: a specific form of
607 programmed cell death? *Experimental Cell Research*. 2003;283(1):1-16.
- 608 45. Bos R, van der Groep P, Greijer AE, Shvarts A, Meijer S, Pinedo HM, et al. Levels of
609 hypoxia-inducible factor-1 α independently predict prognosis in patients with lymph node
610 negative breast carcinoma. *Cancer*. 2003;97(6):1573-81.
- 611 46. Vaupel P, Mayer A. Hypoxia in cancer: significance and impact on clinical outcome.
612 *Cancer and Metastasis Reviews*. 2007;26(2):225-39.
- 613







