

# Multiscale modeling to explore Ebola vaccination strategies

Van Kinh Nguyen<sup>1,2</sup>, Rafael

Mikolajczyk<sup>2,3,4,5</sup> and Esteban A.

Hernandez-Vargas<sup>1,2</sup>

## Subject Areas:

epidemiology, computational biology,  
systems biology

## Keywords:

multiscale, Ebola, epidemics,  
modelling, network model,  
within-host, between-hosts

## Author for correspondence:

Van Kinh Nguyen

e-mail: [knguyen@fias.uni-frankfurt.de](mailto:knguyen@fias.uni-frankfurt.de)

Esteban A. Hernandez-Vargas

e-mail: [vargas@fias.uni-frankfurt.de](mailto:vargas@fias.uni-frankfurt.de)

<sup>1</sup>Frankfurt Institute for Advanced Studies,  
Ruth-Moufang-Strasse 1, 60438, Frankfurt am Main,  
Germany

<sup>2</sup>Helmholtz Centre for Infection Research,  
Braunschweig, Germany

<sup>3</sup>German Centre for Infection Research, Germany

<sup>4</sup>Hannover Medical School, Hannover, Germany

<sup>5</sup>Martin-Luther University, Halle-Wittenberg, Germany

The power to assess potential outcomes and intervention strategies is critical for epidemic preparedness. But emerging and mutated pathogens always challenge our current knowledge, pleading for fresh approaches to explore their epidemic potentials up front.

This paper coupled a within-host viral dynamics model and a between-host network model of Ebola virus (EBOV) infection showing that its transmission characteristics can be faithfully recapitulated.

Based on this multiscale model, EBOV's incubation period is predicted in the range from 2.6 to 12.4 days, while infected subjects can remain infectious until day 17. The predicted basic reproductive number ( $R_0$ ) differs by age-groups: the overall is 1.4 and the highest is 4.7 for the 10-14 years old. Random vaccination strategies can reduce  $R_0$  and case-fatality rate, eliminate the possibility of large outbreaks, but the effect depends on timing and coverage.


A random vaccination program can reduce  $R_0$  below one if 85% coverage is achieved, and if it was conducted during the period from five months before to one week after the start of an epidemic. A vaccination coverage of 33% can reduce the epidemic size by ten to hundred times compared to a non-intervention scenario. Altogether, infection characteristics and epidemic mitigation approaches could be assessed using experimental data. An early, age-group specific, and high coverage vaccination program is the most beneficial.

## 1. Introduction

2

3 Epidemics of infectious diseases are listed among the potential catastrophes and can be potentially  
4 misused as mass destruction weapons [1]. Overwhelming research efforts have been developed  
5 to early predict the danger of the epidemics but their crisis nature left scientists no better option  
6 than learning from the past [1,2]. However, confronting outbreaks of emerging infections requires  
7 swift responses and thus the ability to evaluate quickly and early potential outcomes [1]. As such,  
8 computer simulations of epidemic models undoubtedly hold the potential as the first-aid toolbox  
9 for decision making amid the crisis [1,3].

10 A majority of epidemic modelling studies has exclusively relied on the availability of outbreak  
11 data [4-6]. This approach requires that sufficient incidence data are available; for example, data  
12 at the end of an epidemic or at least until its peak [7]. As such, it has limited applicability to  
13 newly emerging epidemics. Moreover, mechanistic models based on outbreak data are often  
14 oversimplified [8]. For example, the effective transmission probability [6] has been usually  
15 simplified as a single parameter that reflects collective effects of the contact rate with the  
16 infectious, the infectivity of the infectious, and the susceptibility of the susceptible. As a result,  
17 these key processes in the disease transmission are lost, especially the transient nature of the  
18 infection course [9]. In reality, the within-host infection process determines key parameters in



**Figure 1. Schematic presentation of the two infections processes in an outbreak.** At the within-host level, viral replication and immune responses race with each other that eventually determines an individual infectivity, for example, his symptoms and possibly behaviours. At the between hosts level, infected individuals make contact(s) with susceptible individual(s) that eventually lead to a transmission, depending on both the infectivity of the infectious and the susceptibility of the susceptible.

17 the disease transmission [9-13]. In an infected subject, interactions between the viruses and  
18 immune responses shape the viral load dynamics that ultimately defines the incubation period,  
19 the transmission potential, and the recovery rate [11,14]. It is also evident that susceptibility to  
20 infection is not the same for all the susceptible but, among others, it is highly correlated with  
21 a subject's age due to the aging of the immune systems [15,16]. Differences in the within-host  
22 infection profile as well as the susceptibility to infection complicate greatly epidemic models but  
23 at the same time underline their influential roles in determining epidemics features.

24  
25 The interplays between within-host infection and between hosts transmission (Fig. 1) led to  
26 arising attempts connecting the two levels [9,11,17-21], but the approach is still at its infancy [10].  
27 Most of these models are conceptual and theoretical [10], and rely on assumptive or previously  
28 obtained parameter estimates [14,22]. Thus, based on our thorough assessments of modelling  
29 within-host EBOV infection previously [12,13,23], we attempt to simulate EBOV transmission  
30 fitness at the population level using explicitly experimental and epidemiological data.

31 In particular, we embedded a within-host infection model of EBOV infection directly into a  
32 network transmission model at population level to simulate epidemic trajectories. Both the used  
33 models were derived based on empirical data of EBOV infection and human contact networks.  
34 Parameters obtained from simulations were then compared to those estimated based on actual  
35 outbreak data and empirical observations. The results showed that using with-host infection  
36 model not only uncovered faithful estimate of the transmission parameters, but also allowed  
37 the evaluations of realistic vaccination effects. In that capacity, epidemic consequences can be  
38 evaluated ahead of time once within-host viral dynamics are available.

## 39 Material and Methods

40 In an EBOV-infected subject, different immune systems components dynamically evolve in  
41 response to the viral replication dynamic. As a result, a series of events is triggered determining  
42 infection outcomes such as infectious status, symptoms, recovery, or death [24–26]. Therefore,  
43 the EBOV replication dynamics within a host were used in this paper to infer transmission  
44 parameters.

### 45 Within-host model

46 Using viral dynamics and immune responses data within a host, mathematical relations can be  
47 defined to test hypothesized infection mechanisms [12,27]. In this context, non-human primates  
48 (NHPs) are the standard animal model for developing EBOV's therapeutics and vaccines in  
49 humans [28,29] which recently has been used to develop an effective vaccine against EBOV  
50 [30]. Epidemiological and pharmacological studies reported that a viral load higher than  
51  $10^6$  copies/mL [29,31] is associated with a higher mortality rate, whereas observations on  
52 experimental data in NHPs showed that a viral load level higher than  $10^6$  TCID<sub>50</sub> was fatal  
53 [24,25]. Here the viral load dynamics were simulated based on the model as follows [13]:

$$54 \quad \frac{dV}{dt} = r_V V \left(1 - \frac{V}{K_V}\right) \left(\frac{V}{I_n + V}\right) \left(1 - \frac{Ab}{K_{Ab}}\right) \quad (1.1)$$

55 where  $r_V$ ,  $K_V$  and  $I_n$  denote the replication rate, the host's carrying capacity, and a constraint  
56 threshold expressing the lag-phase growth of the virus. The parameter  $K_{Ab}$  represents the  
57 strength of the immune system at which the antibody titre inhibits the viral net growth rate [13].  
58 The model parameters were obtained previously [13] using two experimental datasets on NHPs  
59 [24,25]. The antibody dynamic ( $Ab$ ) was fitted previously [13] to data of NHPs vaccinated with  
60 vesicular stomatitis virus (VSV-EBOV) vaccine [25]. The VSV-EBOV has recently showed efficacy  
61 in human [30]. Detailed of model fitting and data can be found in [13].

### 62 Simulated subject-specific infection course

63 To simulate subject-specific infection course, the antibody response strength  $K_{Ab}$  was varied  
64 from a normal level approximately  $10^{2.5}$  [25,32] to the highest observed level of  $10^{4.5}$  [25].  
65 This value was assumed to vary based on individual's age, i.e., a U-shaped function of age  
66 with larger values for the infant and the elderly [15]. As infective dose can alter the course of  
67 infection [33], the initial condition  $V(0)$  of model Eq. (1.1) was varied depending on from whom a  
68 subject acquires the infection, i.e., equals the lethal dose ( $V_c = 10^{0.15}$  [13]) times the transmission  
69 potential of whom transmits the disease. Here we assumed a direct relation [10] between the  
70 transmission potential and the viral load at the time of infection, i.e., the transmission potential  
71  $p_{\text{Trans}}(t) = V(t)/K_V$ . Note that  $p_{\text{Trans}}(t) = 1$  does not guarantee a successful transmission, but  
72 it was considered collectively with its contacts susceptibility and with the existence of such a  
73 contact.

**Table 1.** Definitions of transmission parameters based on viral load and epidemics outcomes based on network model.

4

Measure	Definition
A Incubation period	the interval between exposure to a pathogen and initial occurrence of symptoms [4] was defined from the infection time to the first time the viral load crosses over the detectable threshold (Fig. S2).
B Time from symptom onset to recovery [4]	defined as the interval between the first day of detectable viral load and the first day the viral load goes undetectable (Fig. S2).
C Time from symptom onset to death [4]	defined as the interval between the first day with detectable viral load and the day the area under the viral load curve (AUC) crosses the reference threshold $AUC_7$ (Fig. S2).
D Basic reproductive number (R0)	calculated based on the network of infected subjects at the end of an epidemic. In terms of network models, this equals the mean degree distribution of the infected network, considering a directed network without loops (e.g., Fig. S3). The R0 by age-group was also calculated in the same fashion based on the assigned age-attribute. Note that in epidemics with intervention, the R0 is called the <i>effective reproductive number</i> (Re).
E Final infected fraction	the proportion of infected nodes at the end of the epidemic simulations.
F Case-fatality rate	the proportion of nodes died as a result of EBOV infection calculated as the end of epidemics.

## 74 Infection outcomes definitions

75 Empirical observations from both EBOV infected human and NHPs showed that the time from  
76 symptom onset to death is approximately one week [24,25,34]. Based on this and the viral load,  
77 we used the total area under the viral load curve (AUC) seven days post-infection in the subjects  
78 that died as a threshold above which the infection is lethal, i.e.,  $AUC_7 = \int_0^7 V(t)dt$ . Otherwise,  
79 infected subjects were assumed recovered once the viral load was no longer detectable (Fig. S2).  
80 Depending on the infective dose and the adaptive immune response strength, an infection will  
81 manifest different viral dynamics. Based on that, we defined the transmission parameters as in  
82 Table 1A-C.

## 83 The network model

84 The European's contact patterns survey data [35] were used to generate a network model  
85 reflecting the number of contacts, the mixing patterns among age-groups, and a specific  
86 population age-structure. The age-distribution of the city Freetown in Sierra Leone was used as  
87 the reference [36]. A detailed description of the implementation can be found in Supplemental 1.  
88 Because EBOV spreads through direct contacts with infectious subjects [33], and that the highest  
89 risk of infection is contacting with blood, faeces, and vomit [37], we used only the data of physical  
90 contacts and excluded those contacts with a duration less than five minutes. To account for the  
91 transmission route through funeral practices in EBOV outbreaks [2], we considered deceased  
92 EBOV-infected subjects infectious until they were buried. During the last epidemics in Sierra  
93 Leone, the time from death to burial was one to two days on average but can be a week [38].  
94 This number was randomly assigned using a truncated normal distribution at zero and seven  
95 with unit mean and variance.

## 96 Transmission outcomes definitions


5

97 To obtain EBOV's epidemics metrics, the within-host infection model was embedded into  
98 network model. Simulations of EBOV epidemic are detailed in Supplemental 2. In short, a  
99 network of ten thousand nodes was generated. Scenarios in which the population was randomly  
100 vaccinated during one-week vaccination programs were tested and compared to a control  
101 simulation without interventions. For each scenario, one thousand simulations were performed,  
102 each of which started with a single random index case. Each time when a contact occurs, the  
103 viral load at the time point was extracted to determine the transmission potential. Next, the  
104 susceptibility of the contact persons were computed as a function of their age [15]. A Bernoulli  
105 trial was then used to determine if the contact results in an infection given the overall transmission  
106 probability. If the transmission succeeds, the newly infected subject has its own infection  
107 profile computed. Based on simulation outputs, the epidemic outcomes were determined as in  
108 Table 1D-F.

## 109 Results

### 110 Basic transmission characteristics

111 Simulations of the outcomes of the within-host infection model showed a highly skewed  
distribution of the basic transmission parameters (Fig. 2). The incubation period derived from




**Figure 2. Extrapolations of the delay distributions post EBOV infection using within-host infection model.** Simulations of the with-host infection model with varied infective dose and immune strengths. The median of the three distributions are 9, 9.7, and 3.8 for A, B, and C, respectively.

112 viral load dynamics ranged from 2.6 to 12.4 days (median: 3.8) compared to the previous estimates  
113 based on actual outbreak data ranging from 3.35 to 12.7 days [4]. The delay time from infection  
114 to recovery ranged from 6.9 to 17.6 days (median: 9.7). Previous estimates of this interval ranged  
115 from 2 to 26 days (median: 10) [4]. The time from infection to death ranged from 8.1 to 15.1 days  
116 (median: 9) compared to previous estimates ranged from 3 to 21 (median: 9–10) [4].  
117

### 118 Basic reproductive number ( $R_0$ )

119 Simulation results showed that the overall estimate of the  $R_0$  was 1.43 (Fig. 3). However, the  
120 estimates differed by age-groups with the highest of 4.7 for the group of 10-14 years of age.  
121 Generally, the age-groups with a higher contact rate had also a higher  $R_0$ . Simulations of  
122 epidemics with varied intervention strategies showed that the  $R_e$  can be reduced below one  
123 if the vaccination program with 85% coverage were deployed as far as five months before the



**Figure 3. Estimates of the basic reproductive number without any intervention, overall and by age-groups.** Simulations of a network of size ten thousand during a period of one year. One thousand simulations were run, each time with a random index case. At the end of each simulation, networks of infected nodes were extracted to compute the average number of secondary infections.

124 introduction of the index case (time zero) or as late as one week after that (Fig. S4). This coverage  
125 threshold was tested as it is the highest vaccine coverage currently achieved worldwide for some  
126 diseases, e.g. Hepatitis B, measles, and polio [39]. Late initiations of similar interventions from  
127 one to five months after the time zero gradually shift the  $R_e$  to the outbreak domain.


128 A lower vaccination coverage of 33% appeared not protective and posed a potential of  
129 outbreak regardless the time of vaccination program (Fig. S4). This coverage was tested as it is a  
130 theoretical protective threshold, i.e.,  $1-1/R_0$  [40]. Note that the tested time window of five months  
131 before the appearance of the index case was chosen based on the windows of opportunity for  
132 EBOV vaccination [13]. As of now, no data are available on the secondary antibody responses to  
133 EBOV; it was assumed that secondary responses are similar to the primary responses.

### 134 Case-fatality rate

135 Simulations showed that the case-fatality rate in the absence of intervention is 90.93% (Fig. S5)  
136 which falls in the range of literature estimates of 0.4 to 0.91 [4]. Furthermore, simulation results  
137 showed that all the intervention strategies mentioned previously can reduce the case-fatality rate.  
138 These results highlight a benefit of vaccination programs even they are late, i.e., they can reduce  
139 the disease severity in newly infected subjects after the vaccination program. As such, relying  
140 solely on  $R_0$  as the indicator for evaluating intervention programs could have overlooked this  
141 life-saving aspect.

### 142 Epidemic final size

143 Theoretical analyses of epidemic models showed when the  $R_0$  is larger than one, the final  
144 size of an epidemic will converge to a two points distribution: either the epidemic dies out  
145 with a small number of infected cases or the epidemic takes off and converges to a normal  
146 distribution [40]. Simulation results confirmed this epidemic behavior (Fig. 4). The results showed  
147 that without intervention, EBOV had approximately 50% to infect more than half the population.  
148 The introduction of vaccination programs at both the coverage thresholds previously mentioned  
149 and at any vaccination time points under assessments were able to scale down the epidemic



**Figure 4. Distribution of the final infected fraction in different timing and coverage of vaccination strategies.** A synthetic population of ten thousand individuals was generated. One thousand simulations were run to simulate the epidemic in the time course of one year. Each time, one individual was chosen randomly as the index case. Circles, diamonds, and connected lines are median. Filled areas are the corresponding non-parametric densities estimates [41]. Two median values are presented for multi-modal density estimates, determining by inflection points.

150 size (Fig. 4). The two points epidemics size distribution gradually converged to a uni-modal  
151 distribution centring at a low infected fraction when the vaccination programs were deployed  
152 earlier. The high vaccine coverage strategy can effectively eliminate the possibilities of having a  
153 major outbreak infecting a large proportion of the population. This can be achieved when the  
154 vaccination programs were deployed any time from a week to five months before time zero.

155 A random vaccination program covering 33% of the population one week before the epidemics  
156 can reduce the final size by more than 100 times compared to a no intervention scenario.  
157 However, the low coverage strategy still showed a small probability that epidemics can become  
158 major whereas the high coverage strategies did not. Vaccination programs deployed during the  
159 epidemics can also substantially reduce the epidemics size. The intervention conducted one  
160 month after time zero can also reduce the final size by more than ten times. These interventions  
161 not only able to reduce the final size, but they can also increase the epidemics extinction  
162 probability.

## 163 Discussion

164 Epidemic modelling aims to obtain generalized solutions to questions such as whether or not a  
165 substantial population fraction is getting infected? how large would the outbreak spread? and  
166 how can the outbreak be mitigated with certain intervention approaches [6,40]. Answering those  
167 questions requires the use of assumptive parameters as well as actual outbreak data [6,14,22,  
168 40]. Our results showed that using information on within-host infection dynamics allows the  
169 identification of those key characteristics in the disease transmission.

170 Estimates of the incubation period suggest a contact tracing period of three weeks for Ebola  
171 epidemics, matching the current WHO's recommendation of 21 days [42]. Estimates of the delay  
172 distributions agreed that EBOV infected subjects can be infectious from day 3 up to three weeks  
173 post infection [4]. Understanding of these delay distributions is critical in both clinical and

174 epidemiological perspectives [43]. These distributions, however, are most often only partially  
175 observed in practice: it is difficult to know the exact time of exposure to the pathogen or  
176 to have complete outbreak data [5,44]. As such, parameter estimation of these distributions  
177 have been relied on testing and comparing different distributional assumptions [44]. In this  
178 paper, mechanistically generated transmission characteristics using viral dynamics remarkably  
179 resemble literature estimates of Ebola. This approach is thus promising and practical given the  
180 accumulating experimental data on varieties of pathogens, notably, the one that as yet unknown  
181 in epidemic contexts.

182 To determine infection outcomes, the threshold  $AUC_7$  was chosen based on suggestions from  
183 empirical data in humans [34] and non-human primates [24,25]. Simulations of the epidemics  
184 using this threshold revealed faithful estimates of the EBOV case-fatality rate (Fig. S5), supporting  
185 the use of the total viral load (AUC) as a criterion for determining infection outcomes. Although  
186 a more precise threshold criterion is desirable, it might not be feasible to obtain in practice  
187 considering inherent ethical reasons. Thus a similar criterion could be considered when adapting  
188 this approach to other infectious diseases, but ideally with dedicated experimental data.

189 Different classes of network models have been proposed, but they cannot reproduce properties  
190 observed in real world networks [45]. Thus, a network model obeying empirical data provides a  
191 more solid ground for epidemic simulations. Apart from mimicking the contact data properties,  
192 our network model can express age-related infection traits via the assigned age attributes. It was  
193 used here to express individual differences in the susceptibility to viral infection - an important  
194 element in a realistic disease transmission. Although contact data might not be available for a  
195 certain target area, the assortative patterns of human contacts and the highly skewed distribution  
196 of the number of contacts might hold true across regions and cultures [35,46]. Thus, the model  
197 demonstrates a simple way to bring empirical data into epidemic modelling studies.

198 Previous  $R_0$  estimates based on EBOV outbreak data were diverse, depending on model  
199 choices and assumptions [4]. Our estimate of  $R_0$  was 1.4 which is within the range of the previous  
200 estimates, ranging from 1.2 to 2.6, with some exceptional estimates up to 4.7 and 8.3 [4]. Notably,  
201 the estimates differed by age-groups with the highest of 4.7 for the group of 10-14-years of age.  
202 Although these estimates depend on Sierra Leon's age-structure, the differences of  $R_0$  estimate  
203 stress the role of the age-structure and contact patterns in the estimation of  $R_0$ , prompting that  
204 age-specific intervention strategies should be considered. The estimates by sub-groups single out  
205 the effort required to control the epidemic [7]. With current assumptions, targeting interventions  
206 to the group 5-20-years of age could be an effective strategy. Note that the differences of  $R_0$  by  
207 age-group could explain the wide variation of the previous estimates where different samples  
208 were employed [4].

209 The following assumptions were used in the paper given the lack of specific experimental data,  
210 but further efforts to produce data are needed to refine and to adapt to other settings: (i) Secondary  
211 antibody responses are the same as primary responses: This underestimates the effect of the  
212 vaccination strategies conducting before the epidemics. Experimental studies on secondary  
213 immune responses to EBOV infection are needed, especially those with a longer follow-up period.  
214 (ii) The transmission potential is directly related to viral load: This is although simple and  
215 reasonable, but different types of relationship, such as non-linear, might exist [10]. Dedicated  
216 animal experiments to define the exact relationship between the viral load the ability to transmit  
217 the virus are needed. (iii) The contact pattern is the same between European countries and  
218 Sierra Leone: Although the contact patterns seemed similar across countries [35], a more sociable  
219 population would increase the estimate of  $R_0$ . (iv) Infection statuses have no influences on the  
220 network structure, except those were buried. This could overestimate  $R_0$  [47]. Taking people's  
221 behaviour changes into epidemic modelling remains a grand challenge [47]. (v) Susceptibility to  
222 EBOV infection is similar to a general viral infection disease: Studies on susceptibility functions  
223 are lacking and require more attentions of the infection research community.

224 Throughout this paper, we showed the possibilities to investigate practical and intriguing  
225 questions using a within-host viral dynamic model. The advantage of this approach is the



226 availability of experimental data and the possibility of conducting experiments to characterize  
227 epidemic transmission. Therefore, *in vitro* and *in vivo* studies of infectious agents could be  
228 seamlessly integrated into studies of between hosts transmission, promoting evidence-based  
229 public health practices.

9

## Funding

This work was supported by the Alfons und Gertrud Kassel-Stiftung and iMed- the Helmholtz Initiative on Personalized Medicine. VKN has been supported by the Presidents Initiative and Networking Funds of the Helmholtz Association of German Research Centres (HGF) under contract number VH-GS-202.

## Authors' contributions

EAHV supervised the project. VKN designed the modelling and performed the simulations. VKN, EAHV and RM provided and analysed the data. VKN, EAHV and RM discussed and wrote the manuscript. All authors reviewed the manuscript.

## Competing interests

The authors declare that they have no any competing interests.

## References

1. B. Gates. The next epidemic—lessons from Ebola. *N. Engl. J. Med.*, 372(15):1381–1384, Apr 2015.
2. P. Piot. Public health: Beating Ebola. *Nature*, 537(7621):484–485, 2016.
3. E. T. Lofgren, M. E. Halloran, C. M. Rivers, et al. Opinion: Mathematical models: a key tool for outbreak response. *Proc. Natl. Acad. Sci. U.S.A.*, 111(51):18095–18096, Dec 2014.
4. M. D. Van Kerkhove, A. I. Bento, H. L. Mills, N. M. Ferguson, and C. A. Donnelly. A review of epidemiological parameters from Ebola outbreaks to inform early public health decision-making. *Sci Data*, 2:150019, 2015.
5. T. Obadia, R. Haneef, and P. Y. Boelle. The R0 package: a toolbox to estimate reproduction numbers for epidemic outbreaks. *BMC Med Inform Decis Mak*, 12:147, Dec 2012.
6. N. Hens, Z. Shkedy, M. Aerts, et al. *Modeling Infectious Disease Parameters Based on Serological and Social Contact Data: A Modern Statistical Perspective*. Statistics for Biology and Health. Springer New York, 2012.
7. J M Heffernan, R J Smith, and L M Wahl. Perspectives on the basic reproductive ratio. *Journal of The Royal Society Interface*, 2(4):281–293, June 2005.
8. N. Mideo, S. Alizon, and T. Day. Linking within- and between-host dynamics in the evolutionary epidemiology of infectious diseases. *Trends Ecol. Evol. (Amst.)*, 23(9):511–517, Sep 2008.
9. S. Alizon, F. Luciani, and R. R. Regoes. Epidemiological and clinical consequences of within-host evolution. *Trends Microbiol.*, 19(1):24–32, Jan 2011.
10. A. Handel and P. Rohani. Crossing the scale from within-host infection dynamics to between-host transmission fitness: a discussion of current assumptions and knowledge. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, 370(1675), Aug 2015.
11. S. C. Chen, C. P. Chio, L. J. Jou, and C. M. Liao. Viral kinetics and exhaled droplet size affect indoor transmission dynamics of influenza infection. *Indoor Air*, 19(5):401–413, Oct 2009.
12. V. K. Nguyen, S. C. Binder, A. Boianelli, M. Meyer-Hermann, and E. A. Hernandez-Vargas. Ebola virus infection modeling and identifiability problems. *Front Microbiol*, 6:257, 2015.
13. V. K. Nguyen and E. A. Hernandez Vargas. Windows of opportunity for Ebola virus infection treatment and vaccination. *bioRxiv*, 2017.
14. S. Lukens, J. DePasse, R. Rosenfeld, et al. A large-scale immuno-epidemiological simulation of influenza A epidemics. *BMC Public Health*, 14:1019, Sep 2014.
15. D. L. Farber, N. A. Yudanin, and N. P. Restifo. Human memory T cells: generation, compartmentalization and homeostasis. *Nat. Rev. Immunol.*, 14(1):24–35, Jan 2014.

16. E. A. Hernandez-Vargas, E. Wilk, L. Canini, et al. Effects of aging on influenza virus infection dynamics. *J. Virol.*, 88(8):4123–4131, Apr 2014.
17. L. N. Murillo, M. S. Murillo, and A. S. Perelson. Towards multiscale modeling of influenza infection. *J. Theor. Biol.*, 332:267–290, Sep 2013.
18. T. Day, S. Alizon, and N. Mideo. Bridging scales in the evolution of infectious disease life histories: theory. *Evolution*, 65(12):3448–3461, Dec 2011.
19. I. M. Longini, A. Nizam, S. Xu, et al. Containing pandemic influenza at the source. *Science*, 309(5737):1083–1087, Aug 2005.
20. N. M. Ferguson, D. A. Cummings, S. Cauchemez, et al. Strategies for containing an emerging influenza pandemic in Southeast Asia. *Nature*, 437(7056):209–214, Sep 2005.
21. X. Cen, Z. Feng, and Y. Zhao. Emerging disease dynamics in a model coupling within-host and between-host systems. *Journal of Theoretical Biology*, 361:141–151, 2014.
22. S. Merler, M. Ajelli, L. Fumanelli, et al. Containing Ebola at the Source with Ring Vaccination. *PLoS Negl Trop Dis*, 10(11):e0005093, Nov 2016.
23. V. K. Nguyen, F. Klawonn, R. Mikolajczyk, and E. A. Hernandez-Vargas. Analysis of Practical Identifiability of a Viral Infection Model. *PLoS ONE*, 11(12):e0167568, 2016.
24. X. Qiu, G. Wong, J. Audet, et al. Reversion of advanced Ebola virus disease in nonhuman primates with ZMapp. *Nature*, 514(7520):47–53, Oct 2014.
25. A. Marzi, S. J. Robertson, E. Haddock, et al. EBOLA VACCINE. VSV-EBOV rapidly protects macaques against infection with the 2014/15 Ebola virus outbreak strain. *Science*, 349(6249):739–742, Aug 2015.
26. J. B. Prescott, A. Marzi, D. Safronetz, et al. Immunobiology of Ebola and Lassa virus infections. *Nat. Rev. Immunol.*, 17(3):195–207, Mar 2017.
27. M. Nowak and R. M. May. *Virus Dynamics : Mathematical Principles of Immunology and Virology: Mathematical Principles of Immunology and Virology*. Oxford University Press, UK, 2000.
28. N. Sullivan, Z. Yang, and G. J. Nabel. Ebola virus pathogenesis: implications for vaccines and therapies. *Journal Of Virology*, 77(18):9733–7, sep 2003.
29. V. Madelain, T. H. Nguyen, A. Olivo, et al. Ebola Virus Infection: Review of the Pharmacokinetic and Pharmacodynamic Properties of Drugs Considered for Testing in Human Efficacy Trials. *Clin Pharmacokinet*, 55(8):907–923, Aug 2016.
30. A. M. Henao-Restrepo, A. Camacho, I. M. Longini, et al. Efficacy and effectiveness of an rVSV-vectored vaccine in preventing Ebola virus disease: final results from the Guinea ring vaccination, open-label, cluster-randomised trial (Ebola AŒa Suffit!). *Lancet*, 389(10068):505–518, Feb 2017.
31. J. Li, H. J. Duan, H. Y. Chen, et al. Age and Ebola viral load correlate with mortality and survival time in 288 Ebola virus disease patients. *Int. J. Infect. Dis.*, 42:34–39, Jan 2016.
32. S. Agarwal and C. Cunningham-Rundles. Assessment and clinical interpretation of reduced IgG values. *Ann. Allergy Asthma Immunol.*, 99(3):281–283, Sep 2007.
33. H. Feldmann and T. W. Geisbert. Ebola haemorrhagic fever. *Lancet (London, England)*, 377(9768):849–62, mar 2011.
34. E. Akerlund, J. Prescott, and L. Tampellini. Shedding of Ebola virus in an asymptomatic pregnant woman. *The New England Journal of Medicine*, 372(25):2467–9, 2015.
35. J. Mossong, N. Hens, M. Jit, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med.*, 5(3):e74, Mar 2008.
36. Statistics Sierra Leone (SSL) and ICF International. *Sierra Leone Demographic and Health Survey 2013*. Freetown, Sierra Leone and Rockville, Maryland, USA: SSL and ICF International., 2014.
37. World Health Organization. *What we know about transmission of the Ebola virus among humans*. Ebola situation assessment, 2014.
38. J. Lipton. Care and burial practices in urban Sierra Leone. Technical report, October 2014.
39. World Health Organization. *Immunization coverage*. Geneva: World Health Organization, 2010.
40. T. Britton. Stochastic epidemic models: A survey. *Mathematical Biosciences*, 225(1):24–35, 2010.
41. A. Bowman and A. Azzalini. *R package sm: nonparametric smoothing methods (version 2.2-5.4)*. University of Glasgow, UK and Università di Padova, Italia, 2014.
42. Centers for Disease Control and Prevention and World Health Organization. *Implementation and management of contact tracing for Ebola virus disease: emergency guideline*, 2015.
43. H. Nishiura. Early efforts in modeling the incubation period of infectious diseases with an acute course of illness. *Emerging Themes in Epidemiology*, 2007.
44. V. Virlogeux, V. J. Fang, M. Park, J. T. Wu, and B. J. Cowling. Comparison of incubation period distribution of human infections with MERS-CoV in South Korea and Saudi Arabia. *Sci Rep*, 6:35839, Oct 2016.

45. R. Pastor-Satorras, C. Castellano, P. Van Mieghem, and A. Vespignani. Epidemic processes in complex networks. *Reviews of Modern Physics*, 87(3):925, 2015.
46. P. Horby, Q. T. Pham, N. Hens, et al. Social contact patterns in Vietnam and implications for the control of infectious diseases. *PLoS ONE*, 6(2):e16965, Feb 2011.
47. S. Funk, S. Bansal, C. T. Bauch, et al. Nine challenges in incorporating the dynamics of behaviour in infectious diseases models. *Epidemics*, 10:21–25, Mar 2015.