

1 **Global & Temporal Patterns of Submicroscopic *Plasmodium falciparum*** 2 **Malaria Infection**

3 **Charles Whittaker¹, Hannah Slater^{1,2}, Teun Bousema³, Chris Drakeley⁴, Azra Ghani¹ and Lucy Okell¹**

4 ¹MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London,
5 London, United Kingdom

6 ²PATH, 2201 Westlake Avenue, Seattle, USA

7 ³Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, The Netherlands

8 ⁴Department of Immunology and Infection, Faculty of Infectious and Tropical Diseases, London School of Hygiene and
9 Tropical Medicine, London, United Kingdom

10

11 **Abstract**

12 Molecular detection of *Plasmodium falciparum* infection has revealed large numbers of individuals
13 with low-density (yet transmissible) infections undetectable by microscopy. Here we present an
14 updated systematic review of cross-sectional malaria surveys to explore the prevalence and drivers of
15 these submicroscopic infections and define where they are likely to be relevant to malaria control
16 efforts. Our results show that submicroscopic infections predominate in low transmission settings, but
17 also reveal marked geographical variation in their prevalence, being highest in South American surveys
18 and lowest in West African studies. Whilst current transmission levels partly explain these results, we
19 find that historical transmission intensity also represents a crucial determinant of the size of the
20 submicroscopic reservoir. Submicroscopic infection was more likely in adults than children, although
21 we did not observe a statistically significant influence of seasonality. Our results suggest that the
22 contribution of submicroscopic infections to transmission likely varies substantially across settings,
23 potentially warranting different approaches to their targeting in the approach to elimination.

24

1 Introduction

2 The ability to accurately detect malaria infection during population surveys in endemic areas is a
3 cornerstone of effective surveillance and control of the parasite. Routinely, this is undertaken using
4 microscopy of blood films or rapid diagnostic tests, although recent years have seen increased usage
5 of more sensitive molecular methods in research contexts. These techniques (typically PCR-based¹)
6 have revealed the widespread presence of infections with parasite densities lower than the threshold
7 of detection by routine methods²⁻⁸.

8 Such “submicroscopic” infections are present across a range of different settings and populations^{5,9-}
9 ¹¹. Although rarely causative of severe symptoms, they have been associated with a number of adverse
10 outcomes during pregnancy^{12,13}, as well as mild anaemia¹⁴ and other symptoms (vomiting, jaundice
11 etc.) in children under 10¹⁵. In addition to their capacity to cause low-grade disease, these infections
12 may hold public health relevance due to their contribution to onwards transmission. Although typically
13 characterised by lower parasite densities than microscopically detectable infections, individuals with
14 submicroscopic infections frequently harbour gametocytes (the transmissible form of the parasite)
15 and are capable of contributing to onwards transmission^{16,17}. Recent work suggests that there is often
16 overlap in the parasite densities (as assessed by quantitative PCR) of individuals that are detectable
17 by microscopy and those who are submicroscopically infected, suggesting that differences in parasite
18 densities (and by extension, infectivity) between these infections may not always be that significant
19 in practice¹⁸. Submicroscopically infected individuals have been shown to contribute to transmission of
20 the parasite across both areas of high¹⁹ and low²⁰ transmission intensity, as well seasonal¹⁶ and
21 perennial¹⁷ settings, further underscoring the potential relevance of this infected sub-group to malaria
22 control efforts.

23 Despite this, our understanding of the factors influencing the size of the submicroscopic reservoir
24 across endemic populations remains far from complete. Previous systematic reviews have found that
25 microscopy misses on average, half of all *P. falciparum* infections compared to PCR-based methods in
26 endemic country cross-sectional surveys²¹ and that adults are more likely to harbour submicroscopic
27 infections than children²². However, these reviews also identified extensive unexplained variation in
28 the size of the submicroscopically infected population across settings, suggesting the existence of
29 other important factors that determine the size of the submicroscopic reservoir. For example,
30 although the burden of submicroscopic infection is highly heterogeneous across different locations^{9,23},
31 it remains unclear whether this represents systematic variation according to geography or is reflective
32 of other underlying location-specific characteristics.

33 Resolving these gaps in our understanding of submicroscopic epidemiology has material
34 consequences for the future of malaria control. Despite mixed reports surrounding more recent
35 progress²⁴, transmission is still declining in many endemic countries and the WHO has identified 21
36 countries that have the potential to eliminate by 2020
37 (<http://www.who.int/malaria/areas/elimination/e2020/en/>). Low transmission settings such as these
38 can possess high proportions of submicroscopically infected individuals^{25,26}: the existence of a
39 significant submicroscopic infection prevalence in an otherwise asymptomatic, microscopically
40 negative population in Ethiopia²⁷ highlights that malaria infections can persist in populations entirely
41 absent of clinical indicators or detectable using typically deployed diagnostics. Understanding the
42 prevalence, detectability and infectiousness of low-density infections in these settings will be essential
43 for planning for elimination: is there benefit in detecting and treating such infections, or are resources

1 better spent elsewhere? Improving our understanding of the drivers of submicroscopic infection in
2 these low transmission areas is therefore crucial to enable better definition of when and where
3 submicroscopic infection is likely to occur and how the size of the submicroscopic reservoir is likely to
4 change as areas approach elimination.

5 Here we update previous reviews on submicroscopic malaria infection prevalence^{21,25}, leveraging the
6 increase in the usage of molecular methods over the past 5 years to explore novel determinants of
7 submicroscopic infection prevalence. These include geographical location and historical patterns of
8 transmission, seasonality and the role of age at a finer resolution than previously possible. These
9 results are then integrated with literature-based estimates of the infectivity of submicroscopic
10 individuals to mosquitoes in order to estimate their contribution to malaria transmission across a
11 range of different settings.

12 **Results**

13 **Microscopic Detectability of Malaria Infection Across the Endemicity Spectrum:** Across the 267
14 surveys included in our analyses that assessed infection status across a range of ages, microscopy
15 detected on average 46.5% of all PCR-detectable infections (43.0% - 50.0%, 95% CI), although this
16 varied substantially across settings (**Fig.1A**). There was no difference in the observed relationship
17 between PCR and microscopy prevalence when the model was fitted separately to data collated in
18 previous reviews and the data newly extracted here (**Supp Fig.2 and Supp Fig.3**): A number of more
19 flexible model structures were also fitted to the data, although a log-linear model provided the best
20 overall fit (**Supp Fig.4**). The overall fitted relationship between PCR prevalence and microscopy
21 prevalence was:

$$22 \quad \text{Log odds PCR prevalence} = 0.94 + 1.16 * \text{Log odds microscopy prevalence}$$

23 where $\log \text{ odds} = \ln(\text{prevalence}/(1-\text{prevalence}))$ and $\text{prevalence} = \exp(\log \text{ odds})/(1+\exp(\log \text{ odds}))$. The
24 prevalence ratio (defined as the proportion of PCR positive infections also detectable by light
25 microscopy) increased as malaria transmission (measured by Survey PCR Prevalence) increased,
26 indicating a declining proportion of submicroscopically infected individuals, from between 70-80% in
27 the areas of lowest PCR prevalence to only 10-20% in the highest prevalence areas (**Fig.1B**). However,
28 there remained large variation in the proportion of submicroscopic infections between surveys carried
29 out in areas with similar PCR prevalence. Grouping surveys by global region (West Africa, East Africa,
30 South America, and Asia/Oceania) revealed marked geographical variation in the prevalence ratio
31 (ANOVA, $p < 0.001$, $df=3$), being lower in South American surveys compared to other regions (Tukey's
32 HSD, $p < 0.001$ for all 3 pairwise comparisons) (**Fig.1C**). There was no statistically significant effect of
33 sampling season after controlling for survey PCR prevalence (ANOVA, $df=2$, $p=0.181$, **Supp Fig.5**), but
34 a statistically significant effect of PCR methodology (ANOVA, $df=5$, $p < 0.001$, **Supp Fig.6**). Scanning a
35 higher number of microscopy fields to determine infection presence/absence was also significantly
36 associated with the prevalence ratio increasing (ANOVA, $df=1$, $p < 0.01$).

37 **Global Variation in the Size of the Submicroscopic Reservoir:** Average transmission intensity (a factor
38 that significantly influences the prevalence ratio, measured here by PCR prevalence) differed
39 significantly between the 4 global regions. It was therefore unclear whether the observed differences
40 in the prevalence ratio were directly related to geography, or whether these results were confounded
41 by the differences in average transmission intensity across the global regions. To account for this, we
42 fitted separate log-linear models to the data from each global region and assessed the modelled

1 prevalence ratio across the range of transmission intensities found in each global region (**Fig.2A-2D**).
2 These results revealed that the prevalence ratio in surveys from South America was lower than would
3 be expected based on their respective transmission intensities alone, and consistently lower than all
4 other global regions (**Fig.2E**). This points to the role of additional factors in shaping the relationship
5 between transmission and microscopy sensitivity other than just endemicity levels. Importantly, there
6 was no evidence that these differences were due to PCR and microscopy methodologies used in
7 different global regions, with no substantial variation in the methodology used between regions (**Supp**
8 **Fig.7**). This highlights that the low prevalence ratio in South America (indicating a wider disparity in
9 PCR and microscopy performance) was likely not occurring due to systematic differences in assay
10 sensitivities between settings.

11 **Exploring the Influence of Historical and Current Transmission Patterns:** The majority of the South
12 American surveys had been undertaken in areas marked by historically low transmission. We
13 therefore investigated whether a high proportion of submicroscopic infections (low prevalence ratio)
14 might be observed in areas around the world that had experienced similar historically low patterns of
15 transmission. Each survey carried out in Africa (a continent for which estimates of malaria prevalence
16 are available at a high spatio-temporal resolution) was geolocated and the estimates of Historical and
17 Current Regional Prevalence at the administrative unit 1 level collated to disentangle the effects of
18 historical prevalence from more contemporary transmission. Note that we distinguish between local
19 malaria transmission (defined by the prevalence recorded in each survey, hereafter referred to as
20 Survey PCR Prevalence), and malaria transmission at the regional level (reflecting broader patterns of
21 transmission). This regional-level transmission represents the average of a heterogeneous mixture of
22 higher and lower transmission areas, and has relevance to local transmission because factors like
23 human movement patterns and circulating parasite genetic diversity are often similar across nearby
24 settings in the same region, even if transmission levels markedly differ^{28,29}.

25 Our results indicated that Regional Historical Prevalence, but not Regional Current Prevalence was a
26 significant predictor of the prevalence ratio when controlling for Survey PCR Prevalence (ANOVA,
27 $p < 0.001$ for both Survey Prevalence and Regional Historical Prevalence, $p = 0.73$ for Regional Current
28 Prevalence), suggesting that historical patterns of transmission are an important determinant of the
29 submicroscopic reservoir size. To further explore this, we classified each survey in our review from
30 Africa into 3 transmission “archetypes” based on historical and current regional levels of transmission
31 and fitted regression models to each. The results from these models were concordant with the results
32 from the ANOVA, with African surveys in regions with both historically and currently low transmission
33 (Sudan, Ethiopia and parts of Kenya and Tanzania) having on average, a lower prevalence ratio (more
34 submicroscopic infections) compared to other low-endemic areas in Africa where historical
35 transmission has been high (**Fig.3B and Fig.3C**). There was no evidence of systematic differences in
36 the PCR and microscopy methodologies in different transmission archetypes (**Supp Fig.8**) and the
37 observed results were robust to the choice of High/Low stratification threshold (**Supp Fig.9**).

38 **The Influence of Age on Submicroscopic Parasite Carriage:** Both increasing age (independent of
39 exposure) and increased immunity (due to previous exposure, which also increases with age) have
40 been linked to lower parasite densities^{30,31}. Previous reviews of submicroscopic epidemiology have
41 confirmed this, although data limitations have previously restricted analyses to only two age
42 categories (≤ 16 and > 16 years old), with a limited number of datapoints in each ($n = 28$ and $n = 13$
43 respectively)²⁵. Motivated by the substantial increase in data generated through the updated
44 systematic review, we defined 3 age-based categories: infants (0-5 years old), older children (5-15


1 years old) and adults (>15 years old), yielding 40, 37 and 43 prevalence survey pairs. The prevalence
2 ratio varied significantly between age groups (ANOVA, $p < 0.001$, $df=2$), and was significantly lower in
3 adults (indicating a greater proportion of submicroscopic infections) compared to young children
4 (Tukey's HSD, $p < 0.001$) and older children (Tukey's HSD, $p < 0.001$). A similar disparity was observed
5 between older and younger children, although this difference was not significant (Tukey's HSD,
6 $p=0.61$). We also explored whether these differences were observed consistently across the range of
7 transmission settings (using survey PCR prevalence as a proxy for transmission intensity) present in
8 the data (**Fig.4A**). Fitting the log-linear regression model separately to the data for each age group
9 highlighted that the increased prevalence ratio observed in infants and older children compared to
10 adults was less pronounced in higher transmission settings (**Fig.4B**). For example, in high endemic
11 areas with 70% overall PCR prevalence, the prevalence ratio for infants was predicted to be 1.42x that
12 of adults, but 1.92x at low endemic areas with 10% overall PCR prevalence. A similar result was
13 observed for adults and older children, suggesting genuine differences in submicroscopic
14 epidemiology both between age groups and across transmission settings.

15 **Contribution of the Submicroscopic Reservoir to Onwards Transmission Across Different Settings:**

16 We next explored how the contribution of submicroscopic infections to onwards transmission might
17 vary across settings characterised by different historical patterns of transmission. To do this, we
18 utilised a range of estimates for the infectivity of submicroscopic infections compared to
19 microscopically detectable infections, as well as the modelled relationship between transmission
20 intensity and the prevalence ratio for each transmission archetype. We estimate that in transmission
21 settings characterised by both historical and current low levels of transmission (the latter defined as
22 $\leq 5\%$ survey prevalence by PCR), submicroscopically infected individuals could account for 17.5% to
23 68.0% of onwards transmission. The exact figure depends on the assumption surrounding the
24 comparative infectivity of microscopically detectable and submicroscopic infections (**Fig.5C**). By
25 contrast, our results suggest the contribution of the submicroscopic reservoir to transmission is less
26 important in settings where transmission has only recently declined (**Fig.5B**) although their
27 contribution is not irrelevant, ranging from 7.8% to 46.0% depending on the assumed comparative
28 infectivity.

29

1
2




3

4 **Figure 1: Prevalence of infection by PCR versus microscopy in 267 prevalence survey pairs and model**
5 **fits.** Bayesian Markov chain Monte Carlo methods were used to fit a linear relationship between PCR
6 prevalence and microscopy prevalence on the log odds scale. **(A)** Microscopy and PCR prevalence (n =
7 267) in surveys identified through this and previous systematic reviews. The fitted model relationship
8 (yellow line) and the 95% credible interval of the mean (paler shaded area). **(B)** The prevalence ratio
9 (the proportion of PCR positive individuals also detectable by microscopy) according to underlying PCR
10 prevalence for each of the 267 survey microscopy-PCR pairs (points) used to fit the full model. The
11 estimated average prevalence ratio (yellow line) and 95% credible interval of the mean (paler shaded
12 area) are also shown. **(C)** Histogram of the prevalence ratio disaggregated by global region. For each
13 region, the size of the point reflects the number of individuals tested by microscopy and PCR. Thick
14 coloured bar on the histogram represents the weighted mean prevalence ratio for each global region.

15

1

2



3

4 **Figure 2: Global variation in the prevalence ratio and the relative size of the submicroscopic**

5 **reservoir.** Microscopy and PCR prevalence in included surveys (points), the model-fitted relationship

6 (coloured line) and 95% credible interval (shaded area) for (A) West Africa (n = 56), (B) East Africa (n


7 = 120), (D) South America (n = 23) and (E) Asia and Oceania (n = 66). (C) The model-fitted average

8 microscopy: PCR prevalence ratio by PCR prevalence for each of the 4 regions (coloured) line and 95%

9 credible interval (shaded area). Coloured countries on each regional map indicates countries for which

10 studies were identified during the systematic review.


1
2
3



4
5
6
7
8
9
10
11
12
13
14
15
16
17

Figure 3: The effect of historical and current transmission intensity on the prevalence of submicroscopic infection in Africa. Historical estimates of malaria prevalence³² were used to assign surveys in Africa to one of three transmission intensity archetypes, and then separate models fitted to each to explore whether variation in the present and historical levels of transmission in a region might influence the frequency of submicroscopic infection in endemic populations. **(A)** Map detailing the African countries for which prevalence surveys were identified, as well as their assigned transmission archetypes based on historical and current transmission intensity (high high= historically high and currently high; high low = historically high, currently low; low low = historically low and currently low) . **(B)** The microscopy: PCR prevalence ratio of surveys in each transmission archetype (dots, $n = 72, 71$ and 26 for high high, high low and low low, respectively), and the modelled average prevalence ratio (coloured line) with 95% credible interval (shaded area). **(C)** The model fitted relationship (coloured lines) and 95% credible interval (shaded areas) for each of the transmission archetypes, with a particular focus on low prevalence (<20%) settings (inset box).


1
2
3
4
5
6
7



8
9
10
11
12
13
14
15
16
17
18

Figure 4: The influence of age on submicroscopic parasite carriage. Separate models were fitted where age disaggregated data were available to assess whether the prevalence of infection by microscopy compared to PCR varied with age. **(A)** Age disaggregated prevalence survey data for young children (0-5 years old, purple points, **n = 40**) older children (5-15 years old, pink points, **n = 37**) and adults (blue points, **n = 43**) with the fitted model relationship (coloured line) and 95% credible interval for each (shaded area). **(B)** The prevalence of microscopy compared to PCR (the prevalence ratio) in surveys where age-disaggregated data (dots) were available by age group, showing the fitted model relationship (coloured lines) and the 95% credible interval (shaded areas).

1
2
3
4



5
6 **Figure 5: The potential contribution of submicroscopic infections to onwads transmission according**
7 **to current and historical transmission intensity.** Using our modelled relationship between PCR and
8 microscopy prevalence (**Fig.3**), we estimated the potential contribution of the submicroscopic
9 reservoir to onwads transmission for each of the transmission archetypes (**A**) High High; (**B**) High Low;
10 (**C**) Low Low) if microscopic infections are either 2x, 5x or 20x more infectious than submicroscopic
11 infections. In each panel, the purple shaded area represents the proportion of transmission
12 attributable to microscopically detectable infections, and the lower shaded areas the contribution of
13 submicroscopic infections for each of the 3 assumed comparative infectivities- darkest shade is 20x
14 (i.e. that microscopically detectable infections are 20x more infectious than submicroscopic
15 infections), intermediate shade is 5x, and lightest shade is 2x.

16

1 Discussion

2 Significant debate surrounds the importance of the submicroscopic reservoir to malaria control efforts
3 and whether they need to be targeted by interventions³³, particularly in areas of low transmission.
4 Disaggregating the now larger quantity of available data (267 prevalence survey pairs from 38
5 countries) has yielded insight into the complex relationships underlying the global pattern of
6 submicroscopic occurrence. In turn, this has facilitated a more refined evaluation of when and where
7 submicroscopic infections are likely to be most prevalent and who is most likely to harbour them. In
8 particular, we show that the transmission gradient in the proportion of submicroscopic infections
9 observed across settings can be largely explained by differences in the historical patterns of
10 transmission and the age structure of the infected population. Moreover, whilst previous work has
11 generally noted the potential relevance of submicroscopic infections in low transmission settings³⁴,
12 our results suggest that this relevance is likely to be highly context dependent, potentially warranting
13 different approaches to their control in different locations.

14 Our results highlight the importance of both individual-level factors (such as age) and setting-specific
15 factors (such as historical patterns of transmission) in determining the size of the submicroscopic
16 reservoir. However, an important caveat to these findings is that there were insufficient data to
17 examine the role of these factors simultaneously. The average age of infection is typically higher in
18 low transmission settings^{35,37}; therefore, a greater proportion of infected individuals in all-age surveys
19 would be expected to be adults. Our results surrounding past transmission history might then be
20 confounded by differences in the average age of infection across these different settings. However,
21 the age distribution of malaria infection appears to adapt fairly rapidly to reflect changes in
22 transmission: in an area of south-western Senegal experiencing a 30-fold reduction in malaria
23 incidence between 1996 and 2010, there was a shift in the age distribution of cases from
24 predominantly under-5s to a distribution that was near uniform across the population³⁸. Similarly,
25 rapid declines in malaria between 2003 and 2007 in The Gambia were shown to be accompanied by
26 an increase in the mean age of paediatric malaria admissions from 3.9 years to 5.6 years³⁹. These
27 clinical results also extend to the prevalence of malaria infection (as assessed by microscopy) in wider
28 populations and communities³⁷. These observations suggest that the difference in infected population
29 age structure between surveys from historically and currently low settings and those from settings
30 where transmission has only recently declined may not be large. Confounding due to differences in
31 age structure are therefore unlikely to explain the observed disparity in the average size of the
32 submicroscopic reservoir across these two types of settings.

33 Another potential limitation is the strong geographical bias in our transmission archetype
34 stratification. The majority of surveys assigned to the “Historically Low, Currently Low” archetype are
35 from East Africa, whilst the majority of surveys in the “Historically High, Currently High” category are
36 from West Africa. The observed results across transmission archetypes could therefore in theory be
37 reflecting geographical variation, rather than variation driven by past transmission history. Two
38 strands of analysis refute this: firstly, that analysis of the submicroscopic reservoir across global
39 regions revealed minimal difference in the prevalence ratio across West and East Africa. And secondly,
40 that the proportion of infections which are submicroscopic in “Historically High, Currently Low”
41 settings (a strata also predominantly composed of studies from East Africa) was consistently lower
42 than that observed for “Historically Low, Currently Low” settings. Together with the results observed
43 for South America, these observed differences support an effect of transmission history on the

1 submicroscopic reservoir which is distinct from the age structure of the infected population and
2 unlikely to be driven by geographical bias.

3 A number of hypotheses could explain these results, including systematic variation in asexual blood
4 stage multiplication rate of *Plasmodium falciparum*⁴⁰ or various haemoglobinopathies and human
5 genetic traits that have been linked to lower average parasite densities^{36,41}. Variation in the extent of
6 parasite genetic diversity in settings characterised by different historical patterns of transmission and
7 the influence of this on the breadth and rate of immunity acquisition might also contribute to the
8 observed results. It is also not possible to definitively preclude systematic variation in diagnostic
9 quality across settings, although this is perhaps unlikely: for example, a recent analysis found that
10 whilst microscopy quality varies across settings, it does not do so systematically with transmission
11 intensity¹⁸. Whilst our results highlight that PCR methodology does significantly influence the
12 prevalence ratio, systematic variation in methodological quality across geographical and transmission
13 archetypes settings was not observed. Interestingly, our analyses revealed no influence of seasonal
14 effects on submicroscopic carriage, despite parasite densities having been shown to slightly rise during
15 the rainy season even when prevalence does not change significantly (possibly due to increased
16 occurrence of superinfection during these periods)^{18,42}. This is consistent with other recent results
17 which similarly found no difference in the submicroscopic reservoir between seasons, when directly
18 comparing the wet and dry season submicroscopic prevalence across the same locations⁴³.

19 Our work suggests that the contribution of submicroscopic infections to onwards transmission is likely
20 to be highly variable across settings. However, this analysis is based on the detection of asexual
21 parasites and does not provide direct insight into gametocyte densities. The relationship between
22 asexual parasite and gametocyte density is highly non-linear³¹ and the distributions of parasite
23 densities in the submicroscopic range can differ substantially between settings¹⁸. The proportion of
24 submicroscopic infections may therefore not linearly relate to their contribution to onwards
25 transmission. For example, whilst a recent membrane feeding study conducted in Burkina Faso and
26 Kenya (high transmission settings) found that 45-75% of all mosquito infections were derived from
27 submicroscopic infections⁴⁷, only 4% of infections arose from submicroscopic individuals in a similar
28 study carried out in Cambodia (a low transmission setting)⁴⁸. These findings contrast with the
29 predictions presented here and underscore the need to better resolve the relationship between
30 submicroscopic parasite carriage, gametocyte densities, and mosquito infectivity.

31 Although progress has stalled recently, substantial enthusiasm still surrounds attempts to eliminate
32 malaria⁴⁹. Despite this however, and despite their potential relevance, our understanding of
33 submicroscopic infections remains far from complete. Namely, do they represent a substantial source
34 of transmission and threat to future progress? Do they need to be targeted in order to achieve malaria
35 elimination? Whilst more work is required, our finding highlight important differences in
36 submicroscopic epidemiology between settings and suggests the absence of a one-size-fits-all solution
37 for malaria control targeting this infected subgroup. Such variation will likely warrant different
38 approaches to malaria control if the infection is to be controlled most effectively in the approach to
39 elimination.

40 **Acknowledgements:** We thank Jenny Walldorf, Miriam Laufer, Andrea Buchwald, Lauren Cohee, Terri
41 Taylor and Don Mathanga for kindly providing access to additional data from their original work.

42 **Contributions:** LO and CD conceived the idea of the study. CW, LO and HS contributed to the design
43 of the study. CW carried out the systematic review and the subsequent analyses, in consultation with

1 LO and HS. CW and LO wrote the manuscript, with HS, TB, CD and AG all providing feedback and
2 suggestions during manuscript drafting. All authors approved the final version of the manuscript.

3 **Competing Interests:** We declare no competing interests

4 **Code and Software:** Complete details of the code used to analyse the collated data can be found at
5 https://github.com/cwhittaker1000/Sub_Patent_Malaria_Analysis

6

Methods

Systematic Review Update and Data Extraction

Malaria prevalence data where both microscopy and PCR based methods had been used to determine infection were compiled, updating a previous review published in 2012²⁵. Searches were carried out using PubMed and Web of Science and the search terms (“PCR” OR “Polymerase Chain Reaction”) AND “falciparum”. Records published January 2010 - July 2017 were searched, yielding 2136 records. A further two records, unpublished at the time of screening, were also included^{20,47}, as were additional data collected through contacting authors of included references. Studies reporting asexual *P.falciparum* prevalence by microscopy and PCR in the same population were included. Surveys of pregnant women, where participants have been chosen on the basis of symptoms/treatment, or did not involve a population from a defined location were excluded. After screening titles and abstracts, 231 references were retained for full text evaluation, of which 60 were included. This yielded 253 new prevalence survey pairs, and 387 total when including previous reviews (which previously identified 45 relevant references)^{21,25} (**Supp Fig.1**). Submicroscopic infections were defined as those where infection was detectable by PCR but not by microscopy. The specificity of microscopy compared to PCR has been shown to be high (average 98.4%²¹); we therefore assume microscopy-positive individuals are also PCR-positive. In the small number of instances where the number of microscopically detected infections was higher than those identified by PCR (n = 9), the prevalence ratio was adjusted to 1 – this adjustment does not qualitatively alter the results described here.

Statistical Analysis and Bayesian Linear Regression

As in previous reviews²², data were analysed using the regression-based methodology detailed in Sharp et al⁵⁰:

$$LM_i = PCR_i + \delta_i \quad (1)$$

$$\delta_i = \delta'_i + \beta_0(PCR_i - \overline{PCR}) \quad (2)$$

where LM_i = log odds of microscopy prevalence in survey i , PCR_i = log odds of PCR prevalence, \overline{PCR} = mean survey PCR prevalence, and δ_i = log odds ratio (OR) of microscopy to PCR prevalence. This formulation allows δ_i to vary between surveys, with β_0 controlling the extent of this variation. This model was fitted within a Bayesian Markov Chain Monte Carlo based framework, implemented in the statistical software package JAGS⁵¹.

Historical and Current Regional Transmission Intensity Stratification

Surveys conducted in African settings were geolocated and prevalence estimates (aggregated to the administrative unit 1 level) from the Malaria Atlas Project⁵² used to characterise current and historical transmission intensity of the region each survey belonged to. We distinguish between local malaria transmission (defined by the prevalence recorded in each survey), and malaria transmission at the regional level (reflecting broader patterns of transmission), with this latter measure used to stratify each study into one of three “transmission archetypes”:

- **Historically High, Currently High:** areas that have historically (defined as 15 years previous to the date of the survey) high transmission intensity (>15% slide prevalence in 2-10 year olds ($PfPR_{2-10}$)) and remain so at the time of the survey (n = 71).
- **Historically High, Currently Low:** Areas of historically high transmission intensity that have declined in recent years to low levels (<15% $PfPR_{2-10}$) (n = 65).

- **Historically Low, Currently Low:** Areas characterised by historical and current low transmission (<15% *PfPR*₂₋₁₀) (n = 28).

Where MAP estimates were unavailable (for dates earlier than 2000), it was assumed that the year 2000 was reflective of historical transmission intensity given the only recent substantial increase in international financing for malaria control (an approximately twentyfold increase between 2000 and 2015)⁵³.

Calculating the Contribution of Submicroscopic Infections to Onwards Transmission

Estimates of comparative infectivity of microscopically-detectable vs submicroscopic infections (the “infectivity ratio”) are variable, ranging from 2x⁵⁴ to a 20x difference⁴⁸. We therefore explored three scenarios where microscopically detectable infections were either 2x, 5x or 20x more infectious to mosquitoes than submicroscopic infections. Proportional contribution to transmission by submicroscopic infections was calculated as:

$$\frac{LM_{-ve}}{(LM_{+ve} * Infectivity Ratio) + LM_{-ve}}$$

The proportional contribution to transmission is the quantity of interest: the relative infectiousness of submicroscopic infections (the prevalence of which is denoted by LM_{-ve}) is therefore set to 1, and *Infectivity Ratio* (2x, 5x or 20x) is a multiplicative factor reflecting the fact that microscopically detectable infections (prevalence denoted by LM_{+ve}) are more infectious. The equation’s denominator reflects total onwards transmission occurring within the population; the numerator the amount of transmission attributable to submicroscopic infections. These analyses assume that that submicroscopic and microscopically infected populations do not differ in other key factors likely to also influence transmission (e.g. age and mosquito exposure).

References

1. Snounou, G. *et al.* The importance of sensitive detection of malaria parasites in the human and insect hosts in epidemiological studies, as shown by the analysis of field samples from Guinea Bissau. *Trans. R. Soc. Trop. Med. Hyg.* **87**, 649–53
2. Lamptey, H. *et al.* The prevalence of submicroscopic Plasmodium falciparum gametocyte carriage and multiplicity of infection in children, pregnant women and adults in a low malaria transmission area in Southern Ghana. *Malar. J.* **17**, 331 (2018).
3. Zhou, Z. *et al.* Assessment of submicroscopic infections and gametocyte carriage of Plasmodium falciparum during peak malaria transmission season in a community-based cross-sectional survey in western Kenya, 2012. *Malar. J.* **15**, (2016).
4. Shekalaghe, S. A. *et al.* Submicroscopic Plasmodium falciparum gametocyte carriage is common in an area of low and seasonal transmission in Tanzania. *Trop. Med. Int. Heal.* **12**, 547–553 (2007).
5. Steenkeste, N. *et al.* Sub-microscopic malaria cases and mixed malaria infection in a remote area of high malaria endemicity in Rattanakiri province, Cambodia: implication for malaria elimination. *Malar. J.* **9**, 108 (2010).
6. Baum, E. *et al.* Common asymptomatic and submicroscopic malaria infections in Western Thailand revealed in longitudinal molecular and serological studies: a challenge to malaria elimination. *Malar. J.* **15**, 333 (2016).
7. Arango, E. M. *et al.* Molecular Detection of Malaria at Delivery Reveals a High Frequency of Submicroscopic Infections and Associated Placental Damage in Pregnant Women from Northwest Colombia. *Am. J. Trop. Med. Hyg.* **89**, 178–183 (2013).
8. Mueller, I. *et al.* High sensitivity detection of Plasmodium species reveals positive correlations between infections of different species, shifts in age distribution and reduced local variation in Papua New Guinea. *Malar. J.* **8**, 41 (2009).
9. Tadesse, F. G. *et al.* The shape of the iceberg: quantification of submicroscopic Plasmodium falciparum and Plasmodium vivax parasitaemia and gametocytaemia in five low endemic settings in Ethiopia. *Malar. J.* **16**, 99 (2017).
10. Rek, J. *et al.* Characterizing microscopic and submicroscopic malaria parasitaemia at three sites with varied transmission intensity in Uganda. *Malar. J.* **15**, 470 (2016).
11. Carmona-Fonseca, J. *et al.* Molecular Detection of Malaria at Delivery Reveals a High Frequency of Submicroscopic Infections and Associated Placental Damage in Pregnant Women from Northwest Colombia. *Am. J. Trop. Med. Hyg.* **89**, 178–183 (2013).
12. Cottrell, G. *et al.* Submicroscopic Plasmodium falciparum Infections Are Associated With Maternal Anemia, Premature Births, and Low Birth Weight. *Clin. Infect. Dis.* **60**, 1481–1488 (2015).
13. Omer, S. A. *et al.* Impact of Submicroscopic Plasmodium falciparum Parasitaemia on Maternal Anaemia and Low Birth Weight in Blue Nile State, Sudan. *J. Trop. Med.* **2019**, 1–7 (2019).
14. Ladeia-Andrade, S., Ferreira, M. U., de Carvalho, M. E., Curado, I. & Coura, J. R. Age-dependent acquisition of protective immunity to malaria in riverine populations of the Amazon Basin of Brazil. *Am. J. Trop. Med. Hyg.* **80**, 452–9 (2009).
15. Katrak, S. *et al.* Clinical consequences of submicroscopic malaria parasitaemia in Uganda. *Malar. J.* **17**, 67 (2018).
16. Ouédraogo, A. L. *et al.* Substantial Contribution of Submicroscopical Plasmodium falciparum Gametocyte Carriage to the Infectious Reservoir in an Area of Seasonal Transmission. *PLoS One* **4**, e8410 (2009).
17. Schneider, P. *et al.* Submicroscopic Plasmodium falciparum gametocyte densities frequently result in mosquito infection. *Am. J. Trop. Med. Hyg.* **76**, 470–4 (2007).
18. Slater, H. *et al.* The density, temporal dynamics and infectiousness of subpatent P. falciparum infections. *Nat. Commun.* (In Press. (2019).
19. Ouédraogo, A. L. *et al.* Dynamics of the Human Infectious Reservoir for Malaria Determined by Mosquito Feeding Assays and Ultra-sensitive Malaria Diagnosis in Burkina Faso. *J. Infect. Dis.* **213**, 90–99 (2016).
20. Tadesse, F. G. *et al.* The Relative Contribution of Symptomatic and Asymptomatic Plasmodium vivax and Plasmodium falciparum Infections to the Infectious Reservoir in a Low-Endemic Setting in Ethiopia. *Clin. Infect. Dis.* (2018). doi:10.1093/cid/cix1123
21. Okell, L. C., Ghani, A. C., Lyons, E. & Drakeley, C. J. Submicroscopic Infection in Plasmodium falciparum –Endemic Populations: A Systematic Review and Meta-Analysis. *J. Infect. Dis.* **200**, 1509–1517 (2009).
22. Okell, L. C. *et al.* Factors determining the occurrence of submicroscopic malaria infections and their relevance for control. *Nat. Commun.* **3**, (2012).
23. Idris, Z. M. *et al.* High and Heterogeneous Prevalence of Asymptomatic and Sub-microscopic Malaria Infections on Islands in Lake Victoria, Kenya. *Sci. Rep.* **6**, (2016).
24. WHO. WHO | World malaria report 2017. WHO (2018).
25. Okell, L. C. *et al.* Factors determining the occurrence of submicroscopic malaria infections and their relevance for control. *Nat. Commun.* **3**, 1237 (2012).
26. Lin, J. T., Saunders, D. L. & Meshnick, S. R. The role of submicroscopic parasitemia in malaria transmission: what is the evidence? *TRENDS Parasitol.* **30**, 183–190 (2014).
27. Tadesse, F. G. *et al.* Submicroscopic carriage of Plasmodium falciparum and Plasmodium vivax in a low endemic area in Ethiopia where no parasitaemia was detected by microscopy or rapid diagnostic test. *Malar. J.* **14**, 303 (2015).
28. Tatem, A. J. & Smith, D. L. International population movements and regional Plasmodium falciparum malaria elimination strategies. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 12222–7 (2010).
29. Tessema, S. *et al.* Using parasite genetic and human mobility data to infer local and cross-border malaria connectivity in Southern Africa. *Elife* **8**, (2019).
30. Rodriguez-Barraquer, I. *et al.* Quantification of anti-parasite and anti-disease immunity to malaria as a function of age and exposure. *Elife* **7**, (2018).
31. Koepfli, C. *et al.* Blood-Stage Parasitaemia and Age Determine Plasmodium falciparum and P-vivax Gametocytaemia in Papua New Guinea. *PLoS One* **10**, (2015).
32. Bhatt, S. *et al.* The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. *Nature* **526**, 207–211 (2015).
33. Lin, J. T., Saunders, D. L. & Meshnick, S. R. The role of submicroscopic parasitemia in malaria transmission: what is the evidence? *Trends Parasitol.* **30**, 183–190 (2014).
34. Cotter, C. *et al.* The changing epidemiology of malaria elimination: new strategies for new challenges. *Lancet* **382**, 900–911 (2013).
35. Carneiro, I. *et al.* Age-patterns of malaria vary with severity, transmission intensity and seasonality in sub-Saharan Africa: a systematic review and pooled analysis. *PLoS One* **5**, e8988 (2010).
36. Drakeley, C., Gonçalves, B., Okell, L. & Slater, H. Understanding the Importance of Asymptomatic and Low-Density Infections for

- Malaria Elimination. in *intechopen.com* (2018).
37. Griffin, J. T., Ferguson, N. M. & Ghani, A. C. Estimates of the changing age-burden of Plasmodium falciparum malaria disease in sub-Saharan Africa. *Nat. Commun.* **5**, 3136 (2014).
 38. Brasseur, P. *et al.* Changing patterns of malaria during 1996-2010 in an area of moderate transmission in Southern Senegal. *Malar. J.* **10**, 203 (2011).
 39. Ceesay, S. J. *et al.* Changes in malaria indices between 1999 and 2007 in The Gambia: a retrospective analysis. *Lancet* **372**, 1545–1554 (2008).
 40. Murray, L. *et al.* Multiplication rate variation in the human malaria parasite Plasmodium falciparum. *Sci. Rep.* **7**, 6436 (2017).
 41. Kiwanuka, G. N. Genetic diversity in Plasmodium falciparum merozoite surface protein 1 and 2 coding genes and its implications in malaria epidemiology: a review of published studies from 1997-2007. *J. Vector Borne Dis.* **46**, 1–12 (2009).
 42. Smith, T. *et al.* Absence of seasonal variation in malaria parasitaemia in an area of intense seasonal transmission. *Acta Trop.* **54**, 55–72 (1993).
 43. Slater, H. C. *et al.* The temporal dynamics and infectiousness of subpatent Plasmodium falciparum infections in relation to parasite density. *Nat. Commun.* **10**, 1433 (2019).
 44. Mosha, J. F. *et al.* Epidemiology of subpatent Plasmodium falciparum infection: implications for detection of hotspots with imperfect diagnostics. *Malar. J.* **12**, (2013).
 45. Roh, M. E. *et al.* High Genetic Diversity of Plasmodium falciparum in the Low-Transmission Setting of the Kingdom of Eswatini. *J. Infect. Dis.* **220**, 1346–1354 (2019).
 46. Bogreau, H. *et al.* Genetic diversity and structure of African Plasmodium falciparum populations in urban and rural areas. *Am. J. Trop. Med. Hyg.* **74**, 953–9 (2006).
 47. Gonçalves, B. P. *et al.* Examining the human infectious reservoir for Plasmodium falciparum malaria in areas of differing transmission intensity. *Nat. Commun.* **8**, 1133 (2017).
 48. Lin, J. T. *et al.* Microscopic Plasmodium falciparum Gametocytemia and Infectivity to Mosquitoes in Cambodia. *J. Infect. Dis.* **213**, 1491–1494 (2016).
 49. Chen, I., Cooney, R., Feachem, R. G. A., Lal, A. & Mpanju-Shumbusho, W. The Lancet Commission on malaria eradication. *Lancet (London, England)* **391**, 1556–1558 (2018).
 50. Sharp, S. J. & Thompson, S. G. Analysing the relationship between treatment effect and underlying risk in meta-analysis: comparison and development of approaches. *Stat. Med.* **19**, 3251–3274 (2000).
 51. Hornik, K., Leisch, F., Zeileis, A. & Plummer, M. *JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs Sampling.*
 52. Bhatt, S. *et al.* The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. *Nature* **526**, 207–211 (2015).
 53. World Health Organisation. WHO | World Malaria Report 2014. (2015).
 54. Coleman, R. E. *et al.* Infectivity of asymptomatic Plasmodium-infected human populations to Anopheles dirus mosquitoes in western Thailand. *J. Med. Entomol.* **41**, 201–8 (2004).